

Product datasheet for TP320641L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NFE4 (NM 001085386) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human transcription factor NF-E4 (NF-E4), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC220641 representing NM_001085386

or AA Sequence: Red=Cloning site Green=Tags(s)

LDPVPRRSAAAIALPRVVCWHTLKSLNGYKNLSSGAETREGLRSSSPVDLPLRPRKQATAAGQRKLLSLQ LLLCACTSVTDLTYWGPAGHGATAPHRSLLAIHLHLVPASSAAMKATGPHNAQTQVNPRGHAPSAEDPTG

SWTVSGPCKDHPHPFLSQSNPPTRISSALPLKTDSALEQTPQQLPSLHLSQG

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK

Predicted MW: 18.9 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001078855

 Locus ID:
 58160

 RefSeq Size:
 967

Cytogenetics: 7q22.1





RefSeq ORF: 576

Synonyms: NF-E4


Summary: The erythroid-specific protein encoded by this gene, and the ubiquitous transcription factor

CP2, form the stage selector protein (SSP) complex, which is involved in preferential

expression of the gamma-globin genes in fetal erythroid cells. Alternate use of an in-frame upstream non-AUG (CUG) translation initiation codon, and a downstream AUG codon, results in two isoforms. While the long isoform (22 kDa) acts as an activator, the short isoform (14 kDa) has been shown to repress gamma-globin gene expression. This gene is located in an

intron of the FBXL13 gene on the opposite strand. [provided by RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified NFE4 protein (Cat# [TP320641]). The protein was produced from HEK293T cells transfected with NFE4 cDNA clone (Cat# [RC220641]) using MegaTran 2.0 (Cat# [TT210002]).