

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP320685

H2AC21 (NM_175065) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins		
Description:	Recombinant protein of human histone cluster 2, H2ab (HIST2H2AB), 20 μg		
Species:	Human		
Expression Host:	HEK293T		
Expression cDNA Clone			
or AA Sequence:	Red=Cloning site Green=Tags(s)		
	MSGRGKQGGKARAKAKSRSSRAGLQFPVGRVHRLLRKGNYAERVGAGAPVYLAAVLEYLTAEILELAGNA ARDNKKTRIIPRHLQLAVRNDEELNKLLGGVTIAQGGVLPNIQAVLLPKKTESHKPGKNK		
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV		
Tag:	C-Myc/DDK		
Predicted MW:	13.8 kDa		
Concentration:	>0.05 µg/µL as determined by microplate BCA method		
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining		
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol		
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.		
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.		
Storage:	Store at -80°C.		
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.		
RefSeq:	<u>NP 778235</u>		
Locus ID:	317772		
UniProt ID:	Q8IUE6		
RefSeq Size:	448		
Cytogenetics:	1q21.2		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	H2AC21 (NM_175065) Human Recombinant Protein – TP320685	
RefSeq ORF:	390	
Synonyms:	H2AB; HIST2H2AB	
Summary:Histones are basic nuclear proteins that are responsible for the nucleosome structure chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of wrapped around a histone octamer composed of pairs of each of the four core histor H2B, H3, and H4). The chromatin fiber is further compacted through the interaction histone, H1, with the DNA between the nucleosomes to form higher order chromati structures. This gene is intronless and encodes a replication-dependent histone tha member of the histone H2A family. Transcripts from this gene contain a palindromi termination element. [provided by RefSeq, Aug 2015]		
Protein Pathway	s: Systemic lupus erythematosus	

Product images:

116 -	-
66 -	-
45 -	_
35 -	-
25 -	-
18 -	
14 -	-

Coomassie blue staining of purified HIST2H2AB protein (Cat# TP320685). The protein was produced from HEK293T cells transfected with HIST2H2AB cDNA clone (Cat# [RC220685]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US