

Product datasheet for TP320900L

KIAA1967 (CCAR2) (NM_021174) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human KIAA1967 (KIAA1967), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC220900 representing NM_021174 Red=Cloning site Green=Tags(s)

MSQFKRQRINPLPGGRNFSGTASTSLLGPPPGLLTPPVATELSQNARHLQGGEKQRVFTGIVTSLHDYFG
 VDDEEVFFQLSVKGRPLQLGKVLVKAAYNPGQAVPWNNAVKVQTLNQLLSPAPPLHVAALGQKQG
 ILGAQPQLIFQPHRIPPLFPQKPLSLFQTSHTLHLSHLNRFARGPHGRDLQGRSDDYDSKRRKQRAGGE
 PWGAKKPRHDLPPYRVHLTPYTVDSPICDFLELQRRYRSLLVPSDFLSVHLSWLSAFPLSQPFLHHP
 IQVSSEKEAAPDAGAEPITADSDPAYSSKVLVLLSSPGLLEELYRCCMLFVDDMAEPRETPHPLKQIKFLL
 GRKEEEAVLVGGWSPSLDGLDPQADPQVLVRTAIRCAQAQTGIDLSGCTKWWRFAEFQYLQPGPPRRRLQ
 TVVVYLPDVTIMPTLEWEALCQQKAAEAPPTQEAQGETEPTQAPDALEQAADTSRRNAETPEATTQ
 QETDLDLPEAPPPLEPAVIARPGCVNLSLHGIVEDRRPKERISFEVMVLAELFLEMLQRDFGYRVYKML
 LSLPEKVVSPPEPEKEEAAKEEATKEEEAIKEEVVKEPKDEAQNENGPATESEAPLKEDGLLPKPLSSGGE
 EEEKPRGEASEDLCEMALDPELLLRDDGEEEFAGAKLEDSEVRSVANSQSEMFEFSSLQDMPKELDPSAV
 LPLDCLLAFVFFDANWCGYLHRRDLERILLTLGIRLSAEQAKQLVSRVVTQNICQYRSLQYSRQEGLDGG
 LPEEVLFGNLDPPLPPGKSTKPGAAPTEHKALVSHNGSLINVGSLQRAEQDQSGRLYLENKIHTLELKL
 EESHNRFSATEVTNKTLAAEMQELRVRLAEAEETARTAERQKSQQLRLLQELRRRLTPLQLEIQRVVEKA
 DSWVEKEEPPASN


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	102.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.



[View online »](#)

Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_066997
Locus ID:	57805
UniProt ID:	Q8N163
RefSeq Size:	4031
Cytogenetics:	8p21.3
RefSeq ORF:	2769
Synonyms:	DBC-1; DBC1; KIAA1967; NET35; p30 DBC; p30DBC
Summary:	<p>Core component of the DBIRD complex, a multiprotein complex that acts at the interface between core mRNP particles and RNA polymerase II (RNAPII) and integrates transcript elongation with the regulation of alternative splicing: the DBIRD complex affects local transcript elongation rates and alternative splicing of a large set of exons embedded in (A + T)-rich DNA regions. Inhibits SIRT1 deacetylase activity leading to increasing levels of p53/TP53 acetylation and p53-mediated apoptosis. Inhibits SUV39H1 methyltransferase activity. As part of a histone H3-specific methyltransferase complex may mediate ligand-dependent transcriptional activation by nuclear hormone receptors. Plays a critical role in maintaining genomic stability and cellular integrity following UV-induced genotoxic stress. Regulates the circadian expression of the core clock components NR1D1 and ARNTL/BMAL1. Enhances the transcriptional repressor activity of NR1D1 through stabilization of NR1D1 protein levels by preventing its ubiquitination and subsequent degradation (PubMed:18235501, PubMed:18235502, PubMed:19131338, PubMed:19218236, PubMed:22446626, PubMed:23352644, PubMed:23398316). Represses the ligand-dependent transcriptional activation function of ESR2 (PubMed:20074560). Acts as a regulator of PCK1 expression and gluconeogenesis by a mechanism that involves, at least in part, both NR1D1 and SIRT1 (PubMed:24415752). Negatively regulates the deacetylase activity of HDAC3 and can alter its subcellular localization (PubMed:21030595). Positively regulates the beta-catenin pathway (canonical Wnt signaling pathway) and is required for MCC-mediated repression of the beta-catenin pathway (PubMed:24824780). Represses ligand-dependent transcriptional activation function of NR1H2 and NR1H3 and inhibits the interaction of SIRT1 with NR1H3 (PubMed:25661920). Plays an important role in tumor suppression through p53/TP53 regulation; stabilizes p53/TP53 by affecting its interaction with ubiquitin ligase MDM2 (PubMed:25732823). Represses the transcriptional activator activity of BRCA1 (PubMed:20160719). Inhibits SIRT1 in a CHEK2 and PSEM3-dependent manner and inhibits the activity of CHEK2 in vitro (PubMed:25361978).[UniProtKB/Swiss-Prot Function]</p>

Product images:

Coomassie blue staining of purified CCAR2 protein (Cat# [TP320900]). The protein was produced from HEK293T cells transfected with CCAR2 cDNA clone (Cat# [RC220900]) using MegaTran 2.0 (Cat# [TT210002]).