

Product datasheet for TP321033

FATP2 (SLC27A2) (NM_003645) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human solute carrier family 27 (fatty acid transporter), member 2 (SLC27A2), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC221033 representing NM_003645 Red=Cloning site Green=Tags(s)

MLSAIYTVLAGLLFLPLLNLCCPYFFQDIGYFLKVAAVGRRVRSYGKRRPARTILRAFLEKARQTPHKP
 FLLFRDETLTYAQVDRRSNQVARALHDHLGLRQGDCVALLMGNEPAYVWLWLGLVKLGCAMAACLNYNI
 RA
 KSL LHCFQCCGAKVLLVSPQLQA AVEEILPSLKKDDVSIYVSRTSNTDGIDSFLDKVDEVSTEPIPESW
 RSEVTFSTPALYIYTS GTTGLPKAAMITHQRIWYGTGLTFVSGLKADDVIYITLPFYHSAALLIGIHGCI
 VAGATLALRTKFSASQFWDDCRKYNVTVIQYIGELRLYLCNSPQKPNDRDHKVR LALGNGLRGD VWRQF
 V
 KRFGDICIEFYAATEGNIGFMNYARKV GAVGRVNYLQKKIITYDLIKYDVEKDEPVRDENG YCVRVPKG
 EVGLLVCKITQLTPFNGYAGAKAQTEKKLRDVFKKGDLYFNSGDLLMVDHENFIYFHDRVGD TFRWKGE
 NVATTEVADTVGLVDFVQEVN VYGVHVPDHEGRIGMASIKMKENHEFDGKKLFQHIADYLP SYARPRFLR
 IQDTIEITGTFKHRKMTLVEEGFNPAVIKDALYFLDDTAKMYVPMTEDIYNAISAKTLKL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	70.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.



[View online »](#)

Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_003636
Locus ID:	11001
UniProt ID:	O14975
RefSeq Size:	2343
Cytogenetics:	15q21.2
RefSeq ORF:	1860
Synonyms:	ACSVL1; FACVL1; FATP2; hFACVL1; HsT17226; VLACS; VLCS
Summary:	The protein encoded by this gene is an isozyme of long-chain fatty-acid-coenzyme A ligase family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. This isozyme activates long-chain, branched-chain and very-long-chain fatty acids containing 22 or more carbons to their CoA derivatives. It is expressed primarily in liver and kidney, and is present in both endoplasmic reticulum and peroxisomes, but not in mitochondria. Its decreased peroxisomal enzyme activity is in part responsible for the biochemical pathology in X-linked adrenoleukodystrophy. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2009]
Protein Families:	Transmembrane
Protein Pathways:	PPAR signaling pathway

Product images:



Coomassie blue staining of purified SLC27A2 protein (Cat# TP321033). The protein was produced from HEK293T cells transfected with SLC27A2 cDNA clone (Cat# [RC221033]) using MegaTran 2.0 (Cat# [TT210002]).