

Product datasheet for TP321915M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BMX (NM 001721) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human BMX non-receptor tyrosine kinase (BMX), transcript variant 2,

100 µg

Species: Human **Expression Host:** HEK293T

Expression cDNA Clone >RC221915 representing NM 001721 or AA Sequence:

Red=Cloning site Green=Tags(s)

MDTKSILEELLLKRSQQKKKMSPNNYKERLFVLTKTNLSYYEYDKMKRGSRKGSIEIKKIRCVEKVNLEE QTPVERQYPFQIVYKDGLLYVYASNEESRSQWLKALQKEIRGNPHLLVKYHSGFFVDGKFLCCQQSCKAA PGCTLWEAYANLHTAVNEEKHRVPTFPDRVLKIPRAVPVLKMDAPSSSTTLAQYDNESKKNYGSQPPSSS TSLAQYDSNSKKIYGSQPNFNMQYIPREDFPDWWQVRKLKSSSSSEDVASSNQKERNVNHTTSKISWEFP ESSSSEEEENLDDYDWFAGNISRSQSEQLLRQKGKEGAFMVRNSSQVGMYTVSLFSKAVNDKKGTVKHYH VHTNAENKLYLAENYCFDSIPKLIHYHQHNSAGMITRLRHPVSTKANKVPDSVSLGNGIWELKREEITLL KELGSGQFGVVQLGKWKGQYDVAVKMIKEGSMSEDEFFQEAQTMMKLSHPKLVKFYGVCSKEYPIYIVTE YISNGCLLNYLRSHGKGLEPSQLLEMCYDVCEGMAFLESHQFIHRDLAARNCLVDRDLCVKVSDFGMTRY VLDDQYVSSVGTKFPVKWSAPEVFHYFKYSSKSDVWAFGILMWEVFSLGKQPYDLYDNSQVVLKVSQGHR

LYRPHLASDTIYQIMYSCWHELPEKRPTFQQLLSSIEPLREKDKH

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW:

77.8 kDa **Concentration:** >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol





Bioactivity: BMX activity verified in a biochemical assay: BMX (BMX non-receptor tyrosine kinase)

(TP321915) activity was measured in a homogeneous time-resolved fluorescent (HTRF \circledR) assay. BMX is a non-receptor tyrosine kinase belonging to the Tec kinase family. BMX has

been implicated in the Stat pathway and reportedly regulates differentiation and

tumorigenicity of several types of cancer cells. Varying concentrations of BMX were added to a reaction mix containing ATP and a biotinylated kinase substrate and the reaction mixture was incubated to allow the protein to phosphorylate the substrate. HTRF detection reagents were then added, and the time-resolved fluorescent signal was measured on a Flexstation 3 microplate reader. The time resolved fluorescent signal is expressed as "delta R" or "\Delta R" and is

a ratio calculated from the fluorescent emission intensities of the donor and acceptor fluors.

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: <u>NP 001712</u>

Locus ID: 660

UniProt ID:P51813RefSeq Size:2514Cytogenetics:Xp22.2

RefSeg ORF: 2025

Synonyms: ETK; PSCTK2; PSCTK3

Summary: This gene encodes a non-receptor tyrosine kinase belonging to the Tec kinase family. The


protein contains a PH-like domain, which mediates membrane targeting by binding to phosphatidylinositol 3,4,5-triphosphate (PIP3), and a SH2 domain that binds to tyrosine-phosphorylated proteins and functions in signal transduction. The protein is implicated in several signal transduction pathways including the Stat pathway, and regulates differentiation and tumorigenicity of several types of cancer cells. Alternatively spliced transcript variants


have been found for this gene. [provided by RefSeq, Mar 2016]

Protein Families: Druggable Genome, Protein Kinase



Product images:





Coomassie blue staining of purified BMX protein (Cat# [TP321915]). The protein was produced from HEK293T cells transfected with BMX cDNA clone (Cat# [RC221915]) using MegaTran 2.0 (Cat# [TT210002]).