

Product datasheet for TP321974L

IL12RB1 (NM_005535) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human interleukin 12 receptor, beta 1 (IL12RB1), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA	>RC221974 representing NM_005535
Clone or AA Sequence:	Red=Cloning site Green=Tags(s)

MEPLVTWVPLLFLFLLSRQGAACRTSECCFQDPPYPDADSGSASGPRDLRCYRISSDRYECQSWQYEGPT
AGVSHFLRCCLSSGRCCYFAAGSATRLQFSDQAGVSVLYTVTLWVESWARNQTEKSPEVTLQLYNSVKYE
PPLGDIKVKLAGQLRMEWETPDNQVGAEVQFRHRTPSSPWKLGDCGPQDDDTESCLCPLMNVAQEFQL
RRRRLGSQGSSWSKWSSPVCVPPENPPQPQVRFVSVEQLGQDGRRLTLKEQPTQLELPEGCQGLAPGTEV
TYRLQLHMLSCPCKAKATRTLHLGKMPYLSGAAYNVAVISSNQFGPGLNQTWHIPADTHTEPVALNISVG
TNGTTMYWPARAQSMYCIWQPVGQDGGLATCSLTAPQDPDPAGMATYSWSRESGAMGQEKCYITIFA
SAHPEKLTWSTVLSTYHFGGNASAAGTPHHVSVKNHLSVSDWAPSLSTCPGVLKEYVVRCRDEDS
KQVSEHPVQPTETQVTL SGLRAGVAYTVQVRADTAWLRGVWSQPQRFSEIVQVSDWLIFASLGSFLSIL
LVGVLGYLGLNRAARHLCPPLPTCASSAIEFPGGKETWQWINPVDFQEEASLQEALVEMSWDKGERTE
PLEKTELPEGAPELALDTELSLEDGDRCKAKM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	70.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_005526
Locus ID:	3594
UniProt ID:	P42701
RefSeq Size:	2100
Cytogenetics:	19p13.11
RefSeq ORF:	1986
Synonyms:	CD212; IL-12R-BETA1; IL12RB; IMD30
Summary:	The protein encoded by this gene is a type I transmembrane protein that belongs to the hemopoietin receptor superfamily. This protein binds to interleukine 12 (IL12) with a low affinity, and is thought to be a part of IL12 receptor complex. This protein forms a disulfide-linked oligomer, which is required for its IL12 binding activity. The coexpression of this and IL12RB2 proteins was shown to lead to the formation of high-affinity IL12 binding sites and reconstitution of IL12 dependent signaling. Mutations in this gene impair the development of interleukin-17-producing T lymphocytes and result in increased susceptibility to mycobacterial and Salmonella infections. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014]
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway

Product images:



Coomassie blue staining of purified IL12RB1 protein (Cat# [TP321974]). The protein was produced from HEK293T cells transfected with IL12RB1 cDNA clone (Cat# [RC221974]) using MegaTran 2.0 (Cat# [TT210002]).