

Product datasheet for TP322239L

DMAP1 (NM_019100) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human DNA methyltransferase 1 associated protein 1 (DMAP1), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC222239 protein sequence Red=Cloning site Green=Tags(s)

MATGADV RDILELGGPEGDAASGTISKKDIINPDKKKSKKSSSETLTFKRPEGMHREVVALLYSDKKDAPP
LLPSDTGQGYRTVKAKLGSKKVRPWKWPFTNPARKDGAMFFHWRRAAEEGKDYPFARFNKTVQVPVYSE
QEYQLYLHDDAWTKAETDHLFDLSRRFDLRFVVIH DRYDHQQFKRSVEDLKERYHICAKLANVRAVPG
TDLKIPVFDAGHERRRKEQLERLYNRTPEQVAEEYLLQELRKIEARKKEREKRSQDLQKLITAADTTAE
QRRTERKAPKKKLPQKKEAEKPAVPETAGIKFPDFKSAGVTLRSQRMKLPSSVGQKKIKALEQM LLELGV
ELSPTPTEELVHMFNELRSDLVLLYELKQACANCEYELQMLRHRHEALARAGVLGGPATPASGPGPASAE
PAVTEPGLGDPDKDTIIDVVGAPLTPNSRKRRESASSSSSVKAKKP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	52.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_061973](#)

Locus ID: 55929

UniProt ID: [Q9NPF5](#)

RefSeq Size: 1784

Cytogenetics: 1p34.1


RefSeq ORF: 1401

Synonyms: DNMAP1; DNMTAP1; EAF2; MEAF2; SWC4

Summary: This gene encodes a subunit of several, distinct complexes involved in the repression or activation of transcription. The encoded protein can independently repress transcription and is targeted to replication foci throughout S phase by interacting directly with the N-terminus of DNA methyltransferase 1. During late S phase, histone deacetylase 2 is added to this complex, providing a means to deacetylate histones in transcriptionally inactive heterochromatin following replication. The encoded protein is also a component of the nucleosome acetyltransferase of H4 complex and interacts with the transcriptional corepressor tumor susceptibility gene 101 and the pro-apoptotic death-associated protein 6, among others. Alternatively spliced transcript variants encoding the same protein have been described. [provided by RefSeq, Jul 2008]

Protein Families: Transcription Factors

Product images:



Coomassie blue staining of purified DMAP1 protein (Cat# [TP322239]). The protein was produced from HEK293T cells transfected with DMAP1 cDNA clone (Cat# [RC222239]) using MegaTran 2.0 (Cat# [TT210002]).