

## Product datasheet for TP322382

### PDI (PDIA2) (NM\_006849) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human protein disulfide isomerase family A, member 2 (PDIA2), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC222382 representing NM_006849 Red=Cloning site Green=Tags(s)

MSRQLLPVLLLLLLLRASCPWGQEQGARSPSEEPPEEEIPKEDGILVLSRHTLGLALREHPALLVEFYAPW  
CGHCQALAPEYSKAAAVLAAESMVVTLAKVDGPAQRELAEEFGVTEYPTLKFFRNGNRTHPEEYTGPRDA  
EGIAEWLRRRVGPSAMRLEDEAAAQALIGGRDLVIGFFQDLQDEDVATFLALAQDALDMTFGLTDRPRL  
FQQFGLTKDVTVLFKFKFDEGRADFPVDEELGLDLGDLRFLVTHSMRLVTEFNSQTSAKIFAARILNHLL  
LFVNQTLAAHRELLAGFGEAAPRFRGQVLFVVDVAADNEHVLQYFGLKAEAAPTLRLVNLETTKKYAPV  
DGGPVTAASITAFCHAVLNGQVKPYLLSQEIPPDWDQRPVKTLVGKNFEQVAFDETKNVFKFYAPWCTH  
CKEMAPAWEALAEKYQDHEDIIIAELDATANELDAFAVHGFPTLKYFPAGPGRKVIEYKSTRDLETFSKF  
LDNGGVLPTEEPPEEPAAPFPEPPANSTMGSKEEL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	58 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online >](#)

RefSeq: [NP\\_006840](#)

Locus ID: 64714

UniProt ID: [Q13087](#)

RefSeq Size: 1726

Cytogenetics: 16p13.3


RefSeq ORF: 1575

Synonyms: PDA2; PDI; PDIP; PDIR

**Summary:** This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, two catalytically active thioredoxin (TRX) domains, two TRX-like domains and a C-terminal ER-retention sequence. The protein plays a role in the folding of nascent proteins in the endoplasmic reticulum by forming disulfide bonds through its thiol isomerase, oxidase, and reductase activity. The encoded protein also possesses estradiol-binding activity and can modulate intracellular estradiol levels. [provided by RefSeq, Sep 2017]

**Protein Families:** Druggable Genome

### Product images:



Coomassie blue staining of purified PDIA2 protein (Cat# TP322382). The protein was produced from HEK293T cells transfected with PDIA2 cDNA clone (Cat# [RC222382]) using MegaTran 2.0 (Cat# [TT210002]).