

Product datasheet for **TP322515M**

ACOX3 (NM_003501) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human acyl-Coenzyme A oxidase 3, pristanoyl (ACOX3), transcript variant 1, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA	>RC222515 representing NM_003501
Clone or AA Sequence:	Red=Cloning site Green=Tags(s)

MASTVEGGDTALLPEFPRGPLDAYRARASFSWKELALFTEGEGMLRFKKTIFSALENDPLFARSPGADLS
LEKYRELNFLRCKRIFEYDFLSVEDMFKSPLKVPALIQCLGMYDSSLAAYLLHSLVFGSAVYSSGSRH
LTYIQKIFRMEIFGCFALTELSHGSNKAIKRTTAHYDPATEEFIIHSPDFEAAKFWVGNMGKTATHAVF
AKLCVPGDQCHGLHPFIVQIRDPKTLMPGVMVGDIGKLLGQNGLDNGFAMFHKVRVPRQSLNRMGDV
TPEGTYVSPFKDVRQRFGASLGSLSGRVSIVSLAILNLKLAVALRFSATRRQFGPTEEEEIPVLEYP
MQQWRLPYLAAYALDHFSKSLFLDLVELQRGLASGDRSARQELGREIHALASASKPLASWTTQQGIQ
ECREACGGHGYLAMNRLGVLRDDNDPNCTYEGDNNILLQQTSNYLLGLLAHQVHDGACFRSPLKSVDFLD
AYPGILDQKFEVSSVADCLDSAVALAAYKWLVCYLLRETYQKLNQEKRSRSSDFEARNKCQVSHGRPLAL
AFVELTVVQRFHEHVHQPSVPPSLRAVLGRLSALYALWLSRHAALLYRGGYFSGEQAGEVLESVAVLALC
SQLKDDAVALVDVIAPPDFVLDSPIGRADGELYKNLWGAVALQESKVLERASWWPEFSVKNPKVIGSLKSKL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	77.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online >](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_003492
Locus ID:	8310
UniProt ID:	O15254
RefSeq Size:	2415
Cytogenetics:	4p16.1
RefSeq ORF:	2100
Summary:	Acyl-Coenzyme A oxidase 3 also know as pristanoyl -CoA oxidase (ACOX3)is involved in the desaturation of 2-methyl branched fatty acids in peroxisomes. Unlike the rat homolog, the human gene is expressed in very low amounts in liver such that its mRNA was undetectable by routine Northern-blot analysis or its product by immunoblotting or by enzyme activity measurements. However the human cDNA encoding a 700 amino acid protein with a peroxisomal targeting C-terminal tripeptide S-K-L was isolated and is thought to be expressed under special conditions such as specific developmental stages or in a tissue specific manner in tissues that have not yet been examined. [provided by RefSeq, Jul 2008]
Protein Pathways:	alpha-Linolenic acid metabolism, Biosynthesis of unsaturated fatty acids, Fatty acid metabolism, Metabolic pathways, PPAR signaling pathway

Product images:



Coomassie blue staining of purified ACOX3 protein (Cat# [TP322515]). The protein was produced from HEK293T cells transfected with ACOX3 cDNA clone (Cat# [RC222515]) using MegaTran 2.0 (Cat# [TT210002]).