

## Product datasheet for **TP322686L**

### ATP6V1G3 (NM\_133326) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human ATPase, H <sup>+</sup> transporting, lysosomal 13kDa, V1 subunit G3 (ATP6V1G3), transcript variant 2, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC222686 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MTSQSQGIHQLLQAEKRAKDKLEEAKKILHLLFLKRRDWDCFWKRKAIEASQGGSGNRRN

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

Tag:	C-Myc/DDK
Predicted MW:	6.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_579872</a>
Locus ID:	127124
UniProt ID:	<a href="#">Q96LB4</a>
RefSeq Size:	691
Cytogenetics:	1q31.3



[View online »](#)


RefSeq ORF: 177

Synonyms: ATP6G3; Vma10

**Summary:** This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'' and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This gene encodes one of three G subunit proteins. Transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

**Protein Pathways:** Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection

### Product images:



Coomassie blue staining of purified ATP6V1G3 protein (Cat# [TP322686]). The protein was produced from HEK293T cells transfected with ATP6V1G3 cDNA clone (Cat# [RC222686]) using MegaTran 2.0 (Cat# [TT210002]).