

Product datasheet for TP322934M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DHPS (NM_013407) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human deoxyhypusine synthase (DHPS), transcript variant 3, 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC222934 representing NM_013407 or AA Sequence: Red=Cloning site Green=Tags(s)

MEGSLEREAPAGALAAVLKHSSTLPPESTQVRGYDFNRGVNYRALLEAFGTTGFQATNFGRAVQQVNAMI EKKLEPLSQDEDQHADLTQSRRPLTSCTIFLGYTSNLISSGIRETIRYLVQHNMVDVLVTTAGGVEEDLI KCLAPTYLGEFSLRGKELRENGINRIGNLLVPNENYCKFEDWLMPILDQMVMEQNTEGVKWTPSKMIARL GKEINNPESVYYWAQKNHIPVFSPALTDGSLGDMIFFHSYKNPGLVLDIVEGARPDEAVSWGKIRVDAQP

VKVYADASLVFPLLVAETFAQKMDAFMHEKNED

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 34.8 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 037539

Locus ID: 1725



 UniProt ID:
 P49366

 RefSeq Size:
 1183

Cytogenetics: 19p13.13

RefSeq ORF: 939

Synonyms: DHS; MIG13


Summary: This gene encodes a protein that is required for the formation of hypusine, a unique amino

acid formed by the posttranslational modification of only one protein, eukaryotic translation initiation factor 5A. The encoded protein catalyzes the first step in hypusine formation by transferring the butylamine moiety of spermidine to a specific lysine residue of the eukaryotic translation initiation factor 5A precursor, forming an intermediate deoxyhypusine

residue. Alternatively spliced transcript variants encoding multiple isoforms have been

observed for this gene. [provided by RefSeq, May 2011]

Product images:



Coomassie blue staining of purified DHPS protein (Cat# [TP322934]). The protein was produced from HEK293T cells transfected with DHPS cDNA clone (Cat# [RC222934]) using MegaTran 2.0

(Cat# [TT210002]).