

## Product datasheet for TP322995L

### Cytochrome P450 2A6 (CYP2A6) (NM\_000762) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human cytochrome P450, family 2, subfamily A, polypeptide 6 (CYP2A6), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC222995 protein sequence Red=Cloning site Green=Tags(s)

MLASGMLLVALLVCLTVMVLMMSVWQQRKSKGKLPFGPTPLPFIGNYLQLNTEQMYNSLMKISERYGPVFT  
IHLGPRRVVLCGHDVREALVDQAEFFSGRGEQATFDWVFKGYGVVFSNGERAKQLRRFSIATLRDFGV  
GKRGIEERIQEEAGFLIDALRGTTGGANIDPTFFLSRTVSNVISSIVFGDRFDYKDKFSLLRMMLGIFQ  
FTSTSTGQLYEMFSSVMKHLPGPQQQAFQLLQGLEDFIAKKVEHNQRTLDPNSPRDFIDSFLIRMQEEEK  
NPNTEFYLNLMVMTTLNLFIGGTETVSTTLRYGFLLLMKHPEVEAKVHEEIDRVIGKNRQPKFEDRAKMP  
YMEAVIHEIQRFQDVIPMSLARRVKKDTKFRDFFLPKGTEVYPMLGSVLRDPSFFSNPQDFNPQHFLNEK  
GQFKKSDAFVFPFSIGKRNCFCGEGLARMELFFFTVMQNFRLKSSQSPKIDVSPKHVGFATIPRNYTMS  
FLPR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	56.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_000753</a>
<b>Locus ID:</b>	1548
<b>UniProt ID:</b>	<a href="#">P11509</a>
<b>RefSeq Size:</b>	1775
<b>Cytogenetics:</b>	19q13.2
<b>RefSeq ORF:</b>	1482
<b>Synonyms:</b>	CPA6; CYP2A; CYP2A3; CYP11A6; P450C2A; P450PB
<b>Summary:</b>	This gene, CYP2A6, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to hydroxylate coumarin, and also metabolizes nicotine, aflatoxin B1, nitrosamines, and some pharmaceuticals. Individuals with certain allelic variants are said to have a poor metabolizer phenotype, meaning they do not efficiently metabolize coumarin or nicotine. This gene is part of a large cluster of cytochrome P450 genes from the CYP2A, CYP2B and CYP2F subfamilies on chromosome 19q. The gene was formerly referred to as CYP2A3; however, it has been renamed CYP2A6. [provided by RefSeq, Jul 2008]
<b>Protein Families:</b>	Druggable Genome, P450, Transmembrane
<b>Protein Pathways:</b>	Caffeine metabolism, Drug metabolism - cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Retinol metabolism

### Product images:



Coomassie blue staining of purified CYP2A6 protein (Cat# [TP322995]). The protein was produced from HEK293T cells transfected with CYP2A6 cDNA clone (Cat# [RC222995]) using MegaTran 2.0 (Cat# [TT210002]).