

## Product datasheet for TP323161

### MATK (NM\_139355) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human megakaryocyte-associated tyrosine kinase (MATK), transcript variant 1, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC223161 representing NM_139355 Red=Cloning site Green=Tags(s)

MAGRGLVSWRAFHGCDSEELPRVSPRFLRAWHPPPVSARMPTRRWAPGTQCITKCEHTRPKPGELAFR  
KGDVVTILEACENKSWYRVKHHTSGQEGLLAAGALREREALSADPKLSLMPWFHGKISGQEAQQLPPE  
DGLFLVRESARHPGDYVLCVSFGRDVIHYRVLHRDGHILTIDEAVFFCNLMDMVEHYSKDKGAICTKLVRP  
KRKHGTKSAEEELARAGWLLNLQHLLTGAQIGEGEFGAVLQGEYLGQKVAVKNIKCDVTAQAFLEDVAVM  
TKMQHENLVRLGVLHQGLYIVMEHVSKGNLVNFLRTRGRALVNTAQLLQFSLHVAEGMEYLESKLLVH  
RDLAARNILVSEDLVAKVSDVDFGLAKAERKGLDSSRLPVKWTAPEALKHGKFTSKSDVWSFGVLLWEVFSY  
GRAPYPKMSLKEVSEAVEKGYRMEPPEGCPGVHVLMSWCWEAEPARRPPFRKLAEKLARELSAGAPAS  
VSGQDADGSTSPRSQEP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	56.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_647612</a>
<b>Locus ID:</b>	4145
<b>UniProt ID:</b>	<a href="#">P42679</a> , <a href="#">F1T0G6</a>
<b>RefSeq Size:</b>	2183
<b>Cytogenetics:</b>	19p13.3
<b>RefSeq ORF:</b>	1521
<b>Synonyms:</b>	CHK; CTK; HHYLTk; HYL; HYLTK; Lsk
<b>Summary:</b>	The protein encoded by this gene has amino acid sequence similarity to Csk tyrosine kinase and has the structural features of the CSK subfamily: SRC homology SH2 and SH3 domains, a catalytic domain, a unique N terminus, lack of myristylation signals, lack of a negative regulatory phosphorylation site, and lack of an autophosphorylation site. This protein is thought to play a significant role in the signal transduction of hematopoietic cells. It is able to phosphorylate and inactivate Src family kinases, and may play an inhibitory role in the control of T-cell proliferation. This protein might be involved in signaling in some cases of breast cancer. Three alternatively spliced transcript variants that encode different isoforms have been described for this gene. [provided by RefSeq, Jul 2008]
<b>Protein Families:</b>	Druggable Genome, Protein Kinase, Stem cell - Pluripotency

### Product images:



Coomassie blue staining of purified MATK protein (Cat# TP323161). The protein was produced from HEK293T cells transfected with MATK cDNA clone (Cat# [RC223161]) using MegaTran 2.0 (Cat# [TT210002]).