

## **Product datasheet for TP323365L**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **UPB1 (NM\_016327) Human Recombinant Protein**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human ureidopropionase, beta (UPB1), 1 mg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC223365 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAGAEWKSLEECLEKHLPLPDLQEVKRVLYGKELRKLDLPREAFEAASREDFELQGYAFEAAEEQLRRPR IVHVGLVQNRIPLPANAPVAEQVSALHRRIKAIVEVAAMCGVNIICFQEAWTMPFAFCTREKLPWTEFAE SAEDGPTTRFCQKLAKNHDMVVVSPILERDSEHGDVLWNTAVVISNSGAVLGKTRKNHIPRVGDFNESTY YMEGNLGHPVFQTQFGRIAVNICYGRHHPLNWLMYSINGAEIIFNPSATIGALSESLWPIEARNAAIANH CFTCAINRVGTEHFPNEFTSGDGKKAHQDFGYFYGSSYVAAPDSSRTPGLSRSRDGLLVAKLDLNLCQQV

NDVWNFKMTGRYEMYARELAEAVKSNYSPTIVKE

**TRTRPL**EQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 43 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

**Storage:** Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 057411

**Locus ID:** 51733





**UniProt ID:** Q9UBR1, A0A024R1H3, B3KNC1

RefSeq Size: 2167

Cytogenetics: 22q11.23 RefSeq ORF: 1152 Synonyms: BUP1

**Summary:** This gene encodes a protein that belongs to the CN hydrolase family. Beta-ureidopropionase

> catalyzes the last step in the pyrimidine degradation pathway. The pyrimidine bases uracil and thymine are degraded via the consecutive action of dihydropyrimidine dehydrogenase (DHPDH), dihydropyrimidinase (DHP) and beta-ureidopropionase (UP) to beta-alanine and


> beta-aminoisobutyric acid, respectively. UP deficiencies are associated with N-carbamyl-betaamino aciduria and may lead to abnormalities in neurological activity. [provided by RefSeq, Jul

2008]

beta-Alanine metabolism, Drug metabolism - other enzymes, Metabolic pathways, **Protein Pathways:** 

Pantothenate and CoA biosynthesis, Pyrimidine metabolism

## **Product images:**



Coomassie blue staining of purified UPB1 protein (Cat# [TP323365]). The protein was produced from HEK293T cells transfected with UPB1 cDNA clone (Cat# [RC223365]) using MegaTran 2.0 (Cat# [TT210002]).