

Product datasheet for TP323397L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cyclophilin F (PPIF) (NM_005729) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human peptidylprolyl isomerase F (PPIF), nuclear gene encoding

mitochondrial protein, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC223397 representing NM_005729

or AA Sequence: Red=Cloning site Green=Tags(s)

MLALRCGSRWLGLLSVPRSVPLRLPAARACSKGSGDPSSSSSSGNPLVYLDVDANGKPLGRVVLELKADV VPKTAENFRALCTGEKGFGYKGSTFHRVIPSFMCQAGDFTNHNGTGGKSIYGSRFPDENFTLKHVGPGVL SMANAGPNTNGSQFFICTIKTDWLDGKHVVFGHVKEGMDVVKKIESFGSKSGRTSKKIVITDCGQLS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 18.8 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 005720

Locus ID: 10105

UniProt ID: P30405, A0A024QZS4



Cyclophilin F (PPIF) (NM_005729) Human Recombinant Protein - TP323397L

RefSeq Size: 2213

Cytogenetics: 10q22.3

RefSeq ORF: 621


Synonyms: Cyp-D; CyP-M; CYP3; CypD

Summary: The protein encoded by this gene is a member of the peptidyl-prolyl cis-trans isomerase

(PPlase) family. PPlases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is part of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of apoptotic and necrotic cell death. [provided by

RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified PPIF protein (Cat# [TP323397]). The protein was produced from HEK293T cells transfected with PPIF cDNA clone (Cat# [RC223397]) using MegaTran 2.0 (Cat# [TT210002]).