

Product datasheet for TP323575M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CTIP1 (BCL11A) (NM_138559) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human B-cell CLL/lymphoma 11A (zinc finger protein) (BCL11A),

transcript variant 3, 100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC223575 representing NM_138559

or AA Sequence: Red=Cloning site Green=Tags(s)

MSRRKQGKPQHLSKREFSPEPLEAILTDDEPDHGPLGAPEGDHDLLTCGQCQMNFPLGDILIFIEHKRKQ CNGSLCLEKAVDKPPSPSPIEMKKASNPVEVGIQVTPEDDDCLSTSSRGICPKQEHIADKLLHWRGLSSP RSAHGALIPTPGMSAEYAPQGICKDEPSSYTCTTCKQPFTSAWFLLQHAQNTHGLRIYLESEHGSPLTPR

VLHTPPFGVVPRELKMCGSFRMEAREPLSSEKI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 26.7 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 612569

Locus ID: 53335





UniProt ID: Q9H165
RefSeq Size: 2358
Cytogenetics: 2p16.1
RefSeq ORF: 729

Synonyms: BCL11A-L; BCL11a-M; BCL11A-S; BCL11A-XL; CTIP1; DILOS; EVI9; HBFQTL5; ZNF856


Summary: This gene encodes a C2H2 type zinc-finger protein by its similarity to the mouse Bcl11a/Evi9

protein. The corresponding mouse gene is a common site of retroviral integration in myeloid leukemia, and may function as a leukemia disease gene, in part, through its interaction with BCL6. During hematopoietic cell differentiation, this gene is down-regulated. It is possibly involved in lymphoma pathogenesis since translocations associated with B-cell malignancies also deregulates its expression. Multiple transcript variants encoding several different

isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Protein Families: Transcription Factors

Product images:



Coomassie blue staining of purified BCL11A protein (Cat# [TP323575]). The protein was produced from HEK293T cells transfected with BCL11A cDNA clone (Cat# [RC223575]) using

MegaTran 2.0 (Cat# [TT210002]).