

## Product datasheet for **TP323681L**

### **CRTAC1 (NM\_018058) Human Recombinant Protein**

#### **Product data:**

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human cartilage acidic protein 1 (CRTAC1), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC223681 representing NM_018058 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MAPSADPGMSRMLPFLLLLWFLPITEGSQRAEPMFTAVTNSVLPPDYDSNPTQLNYGVAVTDVDHDGDF  
E  
IVVAGYNGPNLVLYKYDRAQKRLVNIADVSSPPYALDRQGNAGVTACDIDGDGREEIYFLNTNNAFS  
GVATYTDKLFKFRNNRWEDILSDEVNVARGVASLFAGRSVACVDRKGSGRYSIYANYAYGNVGPDALIE  
MDPEASDLSRGILALRDVAEAGVSKYTGGRGVSVGPILSSASDIFCDNENGNPFLFHNRGDGTVDAA  
ASAGVDDPHQHGRGVALADFNRDGKVDIVYGNWNGPHRLYLQMSTHGKVRFRDIASPKFSMPSPVRTV  
IT  
ADFDNDQELEIFFNNIAYRSSSANRLFRVIRREHGDPLIEELNPGDALEPEGRGTGGVWTFDGDGMLDL  
ILSHGESMAQPLSVFRGNQGFNNNWLVRVPRTRFGAFARGAKVWLYTKKSGAHLRIIDGGSGYLCEMEPV  
AHFGLGKDEASSVEVTWPDGKMVSRNVASGEMNSVLEILYPRDEDTLQDPAPLECGQGFSQQENGHCM  
DT  
NECIQFPFVCPRDKPVCVNTYGSYRCRTNKKCSRGYEPNEDGTACVGTGQSPGPRPTTPTAAAATAAAA  
AAAGAATAAPVLVDGDLNLGSVVKESCEPSC

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**


Tag:	C-Myc/DDK
Predicted MW:	71.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.



[View online »](#)

<b>Note:</b>	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
<b>Storage:</b>	Store at -80°C.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_060528</a>
<b>Locus ID:</b>	55118
<b>UniProt ID:</b>	<a href="#">Q9NQ79</a>
<b>RefSeq Size:</b>	2889
<b>Cytogenetics:</b>	10q24.2
<b>RefSeq ORF:</b>	1983
<b>Synonyms:</b>	ASPIC; ASPIC1; CEP-68; CEP68; LOTUS
<b>Summary:</b>	This gene encodes a glycosylated extracellular matrix protein that is found in the interterritorial matrix of articular deep zone cartilage. This protein is used as a marker to distinguish chondrocytes from osteoblasts and mesenchymal stem cells in culture. The presence of FG-GAP motifs and an RGD integrin-binding motif suggests that this protein may be involved in cell-cell or cell-matrix interactions. Copy number alterations in this gene have been observed in neurofibromatosis type 1-associated glomus tumors. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011]

## Product images:



Coomassie blue staining of purified CRTAC1 protein (Cat# [TP323681]). The protein was produced from HEK293T cells transfected with CRTAC1 cDNA clone (Cat# [RC223681]) using MegaTran 2.0 (Cat# [TT210002]).