

## **Product datasheet for TP323838**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **BPGM (NM 001724) Human Recombinant Protein**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human 2,3-bisphosphoglycerate mutase (BPGM), transcript variant 1,

20 µg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC223838 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSKYKLIMLRHGEGAWNKENRFCSWVDQKLNSEGMEEARNCGKQLKALNFEFDLVFTSVLNRSIHTAW

LI

LEELGQEWVPVESSWRLNERHYGALIGLNREQMALNHGEEQVRLWRRSYNVTPPPIEESHPYYQEIYNDR RYKVCDVPLDQLPRSESLKDVLERLLPYWNERIAPEVLRGKTILISAHGNSSRALLKHLEGISDEDIINI

TLPTGVPILLELDENLRAVGPHQFLGDQEAIQAAIKKVEDQGKVKQAKK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 29.8 kDa

Concentration:  $>0.05 \mu g/\mu L$  as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001715

Locus ID: 669





UniProt ID: P07738

RefSeq Size: 1800 Cytogenetics: 7q33 777 RefSeq ORF:

Synonyms: DPGM; ECYT8

**Summary:** 2,3-diphosphoglycerate (2,3-DPG) is a small molecule found at high concentrations in red


> blood cells where it binds to and decreases the oxygen affinity of hemoglobin. This gene encodes a multifunctional enzyme that catalyzes 2,3-DPG synthesis via its synthetase activity, and 2,3-DPG degradation via its phosphatase activity. The enzyme also has phosphoglycerate phosphomutase activity. Deficiency of this enzyme increases the affinity of cells for oxygen. Mutations in this gene result in hemolytic anemia. Multiple alternatively spliced variants,

encoding the same protein, have been identified. [provided by RefSeq, Sep 2009]

**Protein Families:** Druggable Genome

**Protein Pathways:** Glycolysis / Gluconeogenesis, Metabolic pathways

## **Product images:**



Coomassie blue staining of purified BPGM protein (Cat# TP323838). The protein was produced from HEK293T cells transfected with BPGM cDNA clone (Cat# [RC223838]) using

MegaTran 2.0 (Cat# [TT210002]).