

Product datasheet for TP323889

PADI2 (NM_007365) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human peptidyl arginine deiminase, type II (PADI2), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC223889 representing NM_007365 Red=Cloning site Green=Tags(s)

MLRERTVRLQYGSRVEAVYVLGTYLWTDVYSAAPAGAQTFSCLKHSEHVWVEVRDGEAEVATNGKQRW
L
LSPSTTLRVTMSQASTEASSDKVTVNYDEEGSIPIDQAGLFLTAIEISLDVDADRDGWEKNNPKKASW
TWGPEGQGAILLVNCDRETPWLPKEDCRDEKYSKEDLDMSQMILRTKGPDRLPAGYEIVLYISMDS
KVGVFYVENPFFGQRYIHILGRRKLYHVVKYTGGSSELLFFVEGLCFPDEGFSLVSIHVSLLLEYMAQDI
PLTPIFTDTVIFRIAPWIMTPNILPPVSFVCCMKDNYLFLKEVKNLVEKTNCELKVCFYLNDRGDRWIQ
DEIEFGYIEAPHKGFVVLDSPRDGNLKDFFVKELGPDFGYVTREPLFESVTSLSDFGNLEVSPVTVN
GKTYPLGRILIGSSFPLSGRRMTKVVRDFLKAQQVQAPVELYSDWLTVGHVDEFMSFVPIPGTKKFLLL
MASTSACYKLFREKQKDGHGGEAIMFKGLGGMSSKRITINKILSNESLVQENLYFQRCLDWNDRDILKKELG
LTEQDIIDLPAFKMDEDHRRARAFFPNMVMNIVLKDLDLGIKPFQVVEECCLEMHVRGLLEPLGLECT
FIDDISAYHKFLGEVHCGTNNRRKPFTEFKWVHMVP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	75.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_031391
Locus ID:	11240
UniProt ID:	Q9Y2J8
RefSeq Size:	2348
Cytogenetics:	1p36.13
RefSeq ORF:	1995
Synonyms:	PAD-H19; PAD2; PDI2
Summary:	<p>This gene encodes a member of the peptidyl arginine deiminase family of enzymes, which catalyze the post-translational deimination of proteins by converting arginine residues into citrullines in the presence of calcium ions. The family members have distinct substrate specificities and tissue-specific expression patterns. The type II enzyme is the most widely expressed family member. Known substrates for this enzyme include myelin basic protein in the central nervous system and vimentin in skeletal muscle and macrophages. This enzyme is thought to play a role in the onset and progression of neurodegenerative human disorders, including Alzheimer disease and multiple sclerosis, and it has also been implicated in glaucoma pathogenesis. This gene exists in a cluster with four other paralogous genes. [provided by RefSeq, Jul 2008]</p>

Product images:



Coomassie blue staining of purified PADI2 protein (Cat# TP323889). The protein was produced from HEK293T cells transfected with PADI2 cDNA clone (Cat# [RC223889]) using MegaTran 2.0 (Cat# [TT210002]).