

Product datasheet for TP323988L

MEKK2 (MAP3K2) (NM_006609) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human mitogen-activated protein kinase kinase kinase 2 (MAP3K2), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC223988 representing NM_006609 Red =Cloning site Green =Tags(s)

MDDQQALNSIMQDLAVLHKASRPALSQETRKAKSSSPKKQNDVRVKFEHRGKEKRILQFPRPVKLEDLRS
KAKIAFGQSMDLHYTNNELVIPLTTQDDLKAVELLDRSIHMKSLKILLVINGSTQATNLEPLPSLEDLD
NTVFGAERKKRLSIIGPTSRDRSSPPPGYIPDELHQVARNGSFTSINSEGEFIPESMDQMLDPLSLSSPE
NSGSGSCPSLDSPLDGESYPKSRMPRAQSYPDNHQEFSDYDNPIFEKFGKGGTYPRRYHVSYHHQYNDG
RKTFFRARRTQGTSLRSPVFSPTDHSLSSTSSGSSIFTPEYDDSRIRRRGSDIDNPTLTVMDISPPSRSP
RAPTNWRLGKLLGQGAFGRVYLCYDVDGTRELAVKQVQFDPDSPETSKEVNALECEIQLLNLLHERIVQ
YYGCLRDPQEKLSIFMEYMPGGSIKDQLKAYGALTENVTRKYTRQILEGVHYLHNSMIVHRDIKGANIL
RDSTGNVKLGDFGASKRLQTIKLSGTGMKSVGTPTYWMSPEVISGEGYGRKADIWSVACTVWEMLTEKPP
WAEFEAMAAIFKIATQPTNPKLPPHVSVDYTRDFLKRIFVEAKLRPSADELLRHMFVHYH

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	69.6 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_006600](#)

Locus ID: 10746

UniProt ID: [Q9Y2U5](#), [A0A024RAH0](#)

RefSeq Size: 3336

Cytogenetics: 2q14.3

RefSeq ORF: 1857


Synonyms: MEKK2; MEKK2B

Summary: The protein encoded by this gene is a member of serine/threonine protein kinase family. This kinase preferentially activates other kinases involved in the MAP kinase signaling pathway. This kinase has been shown to directly phosphorylate and activate I κ B kinases, and thus plays a role in NF- κ B signaling pathway. This kinase has also been found to bind and activate protein kinase C-related kinase 2, which suggests its involvement in a regulated signaling process. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Gap junction, GnRH signaling pathway, MAPK signaling pathway

Product images:



Coomassie blue staining of purified MAP3K2 protein (Cat# [TP323988]). The protein was produced from HEK293T cells transfected with MAP3K2 cDNA clone (Cat# [RC223988]) using MegaTran 2.0 (Cat# [TT210002]).