

## Product datasheet for **TP324398L**

### ErbB 3 (ERBB3) (NM\_001005915) Human Recombinant Protein

#### Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human v-erb-b2 erythroblastic leukemia viral oncogene homolog 3 (avian) (ERBB3), transcript variant s, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC224398 representing NM_001005915 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	MRANDALQVLGLLFSLARGSEVGNLSQAVCPGTLNGLSVTGDAENQYQTLKLYERCEWVMGNLEIVLTGH NADLSFLQWIREVTGYVLVAMNEFSTLPLPNLRVVRGTQVYDYGKFAIFVMLNNTNSSHALRQLRLTQLT GQFPMVPSGLTPQQAQDWYLLDDDDPRLLTLSASSKVPVTLAAV
	<b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-Myc/DDK
Predicted MW:	18.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_001005915</a>
Locus ID:	2065
UniProt ID:	<a href="#">P21860</a>



[View online »](#)

RefSeq Size:	1050
Cytogenetics:	12q13.2
RefSeq ORF:	549
Synonyms:	c-erbB-3; c-erbB3; ErbB-3; erbB3-S; FERLK; HER3; LCCS2; MDA-BF-1; p45-sErbB3; p85-sErbB3; p180-ErbB3
Summary:	<p>This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been characterized. One isoform lacks the intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized. [provided by RefSeq, Jul 2008]</p>
Protein Families:	Adult stem cells, Druggable Genome, Protein Kinase, Secreted Protein, Stem cell - Pluripotency, Transmembrane
Protein Pathways:	Calcium signaling pathway, Endocytosis, ErbB signaling pathway

### Product images:



Coomassie blue staining of purified ERBB3 protein (Cat# [TP324398]). The protein was produced from HEK293T cells transfected with ERBB3 cDNA clone (Cat# [RC224398]) using MegaTran 2.0 (Cat# [TT210002]).