

## **Product datasheet for TP324843M**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NAA60 (NM\_001083600) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human N-acetyltransferase 15 (GCN5-related, putative) (NAT15),

transcript variant 3, 100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC224843 representing NM 001083600

or AA Sequence: Red=Cloning site Green=Tags(s)

MTEVVPSSALSEVSLRLLCHDDIDTVKHLCGDWFPIEYPDSWYRDITSNKKFFSLAATYRGAIVGMIVAE IKNRTKIHKEDGDILASNFSVDTQVAYILSLGVVKEFRKHGIGSLLLESLKDHISTTAQDHCKAIYLHVL TTNNTAINFYENRDFKQHHYLPYYYSIRGVLKDGFTYVLYINGGHPPWTILDYIQHLGSALASLSPCSIP

HRVYRQAHSLLCSFLPWSGISSKSGIEYSRTM

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK

**Predicted MW:** 27.3 kDa

Concentration:  $>0.05 \mu g/\mu L$  as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 001077069

**Locus ID:** 79903



ORIGENE

**UniProt ID:** Q9H7X0 2642 RefSeq Size: Cytogenetics: 16p13.3 RefSeq ORF: 726


Synonyms: HAT4; hNaa60; NAT15; NatF

**Summary:** This gene encodes an enzyme that localizes to the Golgi apparatus, where it transfers an

> acetyl group to the N-terminus of free proteins. This enzyme acts on histones, and its activity is important for chromatin assembly and chromosome integrity. Alternative splicing and the use of alternative promoters results in multiple transcript variants. The upstream promoter is located in a differentially methylated region (DMR) and undergoes imprinting; transcript variants originating from this position are expressed from the maternal allele. [provided by

RefSeq, Nov 2015]

## **Product images:**



Coomassie blue staining of purified NAA60 protein (Cat# [TP324843]). The protein was produced from HEK293T cells transfected with NAA60 cDNA clone (Cat# [RC224843]) using

MegaTran 2.0 (Cat# [TT210002]).