

## Product datasheet for **TP324895L**

### **MCG10 (PCBP4) (NM\_033010) Human Recombinant Protein**

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens poly(rC) binding protein 4 (PCBP4), transcript variant 4, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC224895 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MSGSDGGLEEEPELSITLTLRMLMHGKEVGSIIIGKKGETVKRIREQSSARITISEGSCPERITTITGSTA  
AVFHAVSMIAFKLDEDLCAAPANGGNVSRPPVTLRLVIPASQCGSLIGKAGTKIKEIRETTGAQVQVAGD  
LLPNSTERAVTVSGVPDAILCVRQICAVILESPPKGATIPYHPSLSLGTVLLSANQGFSVQGGYQAVTP  
AEVTKLQQLSSHAVPFATPSVWPGLDPGTQTSSQEFLVPNDLIGCVIGRQGSKISEIRQMSGAHIKIGNQ  
AEGAGERHVTITGSPVSIALAQYLITACLETAKSTSGGTPSSAPADLPAPFSPPLTALPTAPPGLLGTPY  
AISLSNFIGLKPMFLALPPASPGPPPGLAAYTAKMAAANGSKKAERQKFSKY

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

Tag:	C-Myc/DDK
Predicted MW:	41.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u><a href="#">NP_127503</a></u>




[View online »](#)

Locus ID:	57060
UniProt ID:	<a href="#">P57723</a> , <a href="#">A0A024R2Y0</a>
RefSeq Size:	1982
Cytogenetics:	3p21.2
RefSeq ORF:	1209
Synonyms:	CBP; LIP4; MCG10

**Summary:** This gene encodes a member of the KH-domain protein subfamily. Proteins of this subfamily, also referred to as alpha-CPs, bind to RNA with a specificity for C-rich pyrimidine regions. Alpha-CPs play important roles in post-transcriptional activities and have different cellular distributions. This gene is induced by the p53 tumor suppressor, and the encoded protein can suppress cell proliferation by inducing apoptosis and cell cycle arrest in G(2)-M. This gene's protein is found in the cytoplasm, yet it lacks the nuclear localization signals found in other subfamily members. Multiple alternatively spliced transcript variants have been described, but the full-length nature for only some has been determined. [provided by RefSeq, Jul 2008]

### Product images:



Coomassie blue staining of purified PCBP4 protein (Cat# [TP324895]). The protein was produced from HEK293T cells transfected with PCBP4 cDNA clone (Cat# [RC224895]) using MegaTran 2.0 (Cat# [TT210002]).