

Product datasheet for TP324975L

Legumain (LGMN) (NM_005606) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human legumain (LGMN), transcript variant 1, 1 mg

Species: Human

Expression Host: HEK293T

**Expression cDNA Clone
or AA Sequence:** >RC224975 protein sequence
Red=Cloning site **Green**=Tags(s)

MWVKVAVFLSVALGIGAIPIDDPEDGGKHWVIVAGSNGWYNYRHQADACHAYQIIHRNGIPDEQIVMM
YDDIAYSEDNPTPGIVINRPNGTDVYQGVPKDYTGEDVTPQNFLAVLRGDAEAVKGGIGSGKVLKSGPQDH
VFIYFTDHGSTGILVFPNEDLHVKDLNETIHYMYKHKMYRKMVFYIEACESGSMNHLDPDNINVYATTA
NPRESSYACYDEKRSTYLGDWYSVNW MEDSDVEDLTKETLHKQYHLVKSHTNTSHVMQYGNKTISTMKV
MQFQGMKRKASSPVPLPPVTHLDLTPSPDVPLTIMKRKLMNTNDLEESRQLTEEIQRHLDARHLIEKSVR
KIVSLLAASEAEVEQLLSERAPLTGHSCYPEALLHFRTHCFNWHSPTYEYALRHLYVLVNLCEKPYPLHR
IKLSMDHVCLGHY

SGPTRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 47.6 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_005597](#)



[View online >](#)


Locus ID: 5641
UniProt ID: [Q99538](#), [Q53XC6](#), [Q96CY7](#)
RefSeq Size: 2073
Cytogenetics: 14q32.12
RefSeq ORF: 1299
Synonyms: AEP; LGMN1; PRSC1

Summary: This gene encodes a cysteine protease that has a strict specificity for hydrolysis of asparaginyl bonds. This enzyme may be involved in the processing of bacterial peptides and endogenous proteins for MHC class II presentation in the lysosomal/endosomal systems. Enzyme activation is triggered by acidic pH and appears to be autocatalytic. Protein expression occurs after monocytes differentiate into dendritic cells. A fully mature, active enzyme is produced following lipopolysaccharide expression in mature dendritic cells. Overexpression of this gene may be associated with the majority of solid tumor types. This gene has a pseudogene on chromosome 13. Several alternatively spliced transcript variants have been described, but the biological validity of only two has been determined. These two variants encode the same isoform. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Protease

Protein Pathways: Antigen processing and presentation, Lysosome

Product images:



Coomassie blue staining of purified LGMN protein (Cat# [TP324975]). The protein was produced from HEK293T cells transfected with LGMN cDNA clone (Cat# [RC224975]) using MegaTran 2.0 (Cat# [TT210002]).