

Product datasheet for TP325258

NACA (NM_001113201) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins Description: Purified recombinant protein of Homo sapiens nascent polypeptide-associated complex alpha subunit (NACA), transcript variant 2, 20 µg Species: Human **Expression Host:** HEK293T **Expression cDNA Clone** >RC225258 representing NM 001113201 or AA Sequence: Red=Cloning site Green=Tags(s) MPGEATETVPATEQELPQPQAETGSGTESDSDESVPELEEQDSTQATTQQAQLAAAAEIDEEPVSKAKQS RSEKKARKAMSKLGLRQVTGVTRVTIRKSKNILFVITKPDVYKSPASDTYIVFGEAKIEDLSQQAQLAAA EKFKVQGEAVSNIQENTQTPTVQEESEEEEVDETGVEVKDIELVMSQANVSRAKAVRALKNNSNDIVNAI MELTM **TRTRPLEQKLISEEDLAANDILDYKDDDDKV** C-Myc/DDK Tag: Predicted MW: 23.2 kDa **Concentration:** >0.05 µg/µL as determined by microplate BCA method **Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol Recombinant protein was captured through anti-DDK affinity column followed by **Preparation:** conventional chromatography steps. Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. Store at -80°C. Storage: Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. RefSeq: NP 001106672 Locus ID: 4666




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	NACA (NM_001113201) Human Recombinant Protein – TP325258
UniProt ID:	<u>Q13765, A0A024RB41</u>
Cytogenetics:	12q13.3
RefSeq ORF:	645
Synonyms:	HSD48; NAC-alpha; NACA1; skNAC
Summary:	This gene encodes a protein that associates with basic transcription factor 3 (BTF3) to form the nascent polypeptide-associated complex (NAC). This complex binds to nascent proteins that lack a signal peptide motif as they emerge from the ribosome, blocking interaction with the signal recognition particle (SRP) and preventing mistranslocation to the endoplasmic reticulum. This protein is an IgE autoantigen in atopic dermatitis patients. Alternative splicing results in multiple transcript variants, but the full length nature of some of these variants, including those encoding very large proteins, has not been determined. There are multiple pseudogenes of this gene on different chromosomes. [provided by RefSeq, Feb 2016]
Protein Families	: Druggable Genome, Transcription Factors

Product images:



Coomassie blue staining of purified NACA protein (Cat# TP325258). The protein was produced from HEK293T cells transfected with NACA cDNA clone (Cat# [RC225258]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US