

Product datasheet for TP325687L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LRRFIP2 (NM 001134369) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Homo sapiens leucine rich repeat (in FLII) interacting protein

2 (LRRFIP2), transcript variant 3, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC225687 representing NM_001134369

or AA Sequence: Red=Cloning site Green=Tags(s)

MGTPASGRKRTPVKDRFSAEDEALSNIAREAEARLAAKRAARAEARDIRMRELERQQKELDEKSDKQYAE NYTRPSSRNSASATTPLSGNSSRRGSGDTSSLIDPDTSLSELRDIYDLKDQIQDVEGRYMQGLKELKESL SEVEEKYKKAMVSNAQLDNEKNNLIYQVDTLKDVIEEQEEQMAEFYRENEEKSKELERQKHMCSVLQHKM EELKEGLRQRDELIEKHGLVIIPDGTPNGDVSHEPVAGAITVVSQEAAQVLESAGEGPLDVRLRKLAGEK EELLSQIRKLKLQLEEERQKCSRNDGTVGDLAGLQNGSDLQFIEMQRDANRQISEYKFKLSKAEQDITTL EQSISRLEGQVLRYKTAAENAEKVEDELKAEKRKLQRELRTALDKIEEMEMTNSHLAKRLEKMKANRTAL

LAQQ

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK
Predicted MW: 48.1 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





RefSeq: NP 001127841

Locus ID: 9209

UniProt ID: Q9Y608, A8K649


Cytogenetics: 3p22.2 RefSeq ORF: 1272 Synonyms: HUFI-2

Summary: The protein encoded by this gene, along with MYD88, binds to the cytosolic tail of toll-like

receptor 4 (TLR4), which results in activation of nuclear factor kappa B signaling. The ubiquitin-like protein FAT10 prevents the interaction of the encoded protein and TLR4, thereby inactivating the nuclear factor kappa B signaling pathway. In addition, this protein can downregulate the NLRP3 inflammasome by recruiting the caspase-1 inhibitor Flightless-I to

the inflammasome complex. [provided by RefSeq, Jan 2017]

Product images:



Coomassie blue staining of purified LRRFIP2 protein (Cat# [TP325687]). The protein was produced from HEK293T cells transfected with LRRFIP2 cDNA clone (Cat# [RC225687]) using MegaTran 2.0 (Cat# [TT210002]).