

## Product datasheet for TP325807L

### Chk1 (CHEK1) (NM\_001114122) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens CHK1 checkpoint homolog ( <i>S. pombe</i> ) (CHEK1), transcript variant 3, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC225807 protein sequence Red=Cloning site Green=Tags(s)

MAVPFVEDWDLVQTLGEGAYGEVQLAVNRVTEEAVAVKIVDMKRAVDCPENIKKEICINKMLNHENVVKF  
YGHRRREGNIQYLFLEYCSGGELFDRIEPDIGMPEPDAQRFFHQLMAGVVYLHGIGITHRDIKPENLLLDE  
RDNLKISDFGLATVFRYNNRERLLNKMCGTLPYVAPELLKRREFHAEPVDVWSCGVILTAMLGELPWDQ  
PSDSCQEYSDWKEKTYLNPWKKIDSAPLALLHKILVENPSARITIPDIKKDRWYNKPLKKGAKRPRVTS  
GGVSESPSGFSKHIQSNLDFSPVNSASSEENVKYSSSQPEPRTGLSLWDTSPSYIDKLVQGISFSQPTCP  
DHMLLNSQLLGTGSSQNPWQRLVKRMTRFFTCLDADKSYQCLKETCEKLGQWKKSCMNQVTISTTDRR  
NNKLIFKVNLLMDDKILVDFRLSKGDGLEFKRHFLKIKGKLIDIVSSQKWLPAT

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	54.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP\\_001107594](#)

Locus ID: 1111

UniProt ID: [O14757](#), [B4DT73](#)

RefSeq Size: 4174

Cytogenetics: 11q24.2

RefSeq ORF: 1428


Synonyms: CHK1

**Summary:** The protein encoded by this gene belongs to the Ser/Thr protein kinase family. It is required for checkpoint mediated cell cycle arrest in response to DNA damage or the presence of unreplicated DNA. This protein acts to integrate signals from ATM and ATR, two cell cycle proteins involved in DNA damage responses, that also associate with chromatin in meiotic prophase I. Phosphorylation of CDC25A protein phosphatase by this protein is required for cells to delay cell cycle progression in response to double-strand DNA breaks. Several alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Oct 2011]

**Protein Families:** Druggable Genome, Protein Kinase, Stem cell - Pluripotency

**Protein Pathways:** Cell cycle, p53 signaling pathway

### Product images:



Coomassie blue staining of purified CHEK1 protein (Cat# [TP325807]). The protein was produced from HEK293T cells transfected with CHEK1 cDNA clone (Cat# [RC225807]) using MegaTran 2.0 (Cat# [TT210002]).