

Product datasheet for TP325964

WEE2 (NM_001105558) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human WEE1 homolog 2 (<i>S. pombe</i>) (WEE2), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC225964 representing NM_001105558 Red =Cloning site Green =Tags(s)
	<p>MDDKDIDKELRQKLNFSYCEETEIEGQKKVEESREASSQTPEKGEVQDSEAKGTPPWTPLSNVHELDTSS EKDKESPDQILRTPVSHPLKCPETPAQPDSRSKLLPSDSPSTPKTMLSRLVISPTGKLPSRGPKHKLTP APLKDEMTSLALVNINPFTPESYKKLFLQSGGKRKIRGDLEEAGPEEGKGGPLAKRCVLRETNMASRYEK EFLEVEKIGVGEFGTVYKCIKRLDGCVYAIKRSMKTFTELSNENSALHEVYAHAVLGHHPHVRYSSWA EDDHMIQNEYCNGGSLQAAISENTKSGNHFEFPKLDILLQISLGLNYIHNSMVLHDIKPSNIFICHK MQSESSGVIEEVENEADWFLSANVMYKIGDLGHATSINKPKVEEGDSRFLANEILQEDYRHLPKADIFAL GLTIAVAAGAESLPTNGAAWHHIRKGNFPDVPQELSEFSLLKNMIQPDAAEQRPSAAAALARNTVLRPSL GKTEELQQQLNLEKFKTATLERELREAQQAQSPQGYTHHGDTGVS GHTGSRSTKRLVGGKSARSSSFTS GEREPLH</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	62.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001099028
Locus ID:	494551
UniProt ID:	P0C1S8
Cytogenetics:	7q34
RefSeq ORF:	1701
Synonyms:	OOMD5; WEE1B
Summary:	Oocyte-specific protein tyrosine kinase that phosphorylates and inhibits CDK1/CDC2 and acts as a key regulator of meiosis during both prophase I and metaphase II (PubMed:29606300). Required to maintain meiotic arrest in oocytes during the germinal vesicle (GV) stage, a long period of quiescence at dictyate prophase I, by phosphorylating CDK1 at 'Tyr-15', leading to inhibit CDK1 activity and prevent meiotic reentry. Also required for metaphase II exit during egg activation by phosphorylating CDK1 at 'Tyr-15', to ensure exit from meiosis in oocytes and promote pronuclear formation (By similarity).[UniProtKB/Swiss-Prot Function]
Protein Pathways:	Cell cycle

Product images:



Coomassie blue staining of purified WEE2 protein (Cat# TP325964). The protein was produced from HEK293T cells transfected with WEE2 cDNA clone (Cat# [RC225964]) using MegaTran 2.0 (Cat# [TT210002]).