

## Product datasheet for TP326803

### NFAT1 (NFATC2) (NM\_001136021) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2 (NFATC2), transcript variant D, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC226803 representing NM_001136021 Red=Cloning site Green=Tags(s)

MQREAAFRLGHCHPLRIMGSVDQEEPNAHKVASPPSGPAYPDDVLDYGLKPYSPLASLSGEPGRFGEPD  
RVGPQKFLSAAKPAGASGLSPRIEITPSHELIQAVGPLRMRDAGLLVEQPPLAGVAASPRFTLPVPGFEG  
YREPLCLSPASSGSSASFISDTFSPYTPSPVSPNNGGPDLLCPQFQNIHAYSPRTSPIMSPRTSLAEDS  
CLGRHSPVPRPASRSSSPGAKRRHSCEALVALPPGASPQRSRSPSPQPSSHVAPQDHGSPAGYPPVAGS  
AVIMDALNSLATDSPCGIPPKMWKTSPPDPSVSAAPSKAGLPRHIYPAVEFLGPCEQGERRNSAPESILL  
VPPTWPKPLVPAIPICSIPTASLPLEWPLSSQSGSYELRIEVQPKPHHRAHYETEGSRGAVKAPTGGH  
PVVQLHGYMENKPLGLQIFIGTADERILKPHAFYQVHRITGKTVTTTSYEKIVGNTKLEIPLPKNNMR  
ATIDCAGILKLRNADIELRKGETDIGRKNTRVRLVFRVHIPESGRIVSLQTASNPIECSQRSHELPMV  
ERQDTSCLVYGGQQMILTGQNFSESKVFTEKTTDGGQIWEWMEATVDKDKSQPNMLFVEIPEYRNKHI  
RTPVKVNFYVINGKRKRSQPQHFTYHPVPAIKTEPTDEYDPTLICSPHGGGLSQPYYPQHPMVAESPSC  
LVATMAPCQQFRTGLSSPDARYQQQNPAAVLYQRSKSLSPSLLGYQQPALMAAPLSLADAHRSVLVHAGS  
QGQSSALLHPSPTNQQASPVHYSPTNQQLRCSGSHQEFQHIMYCENFAPGTRPGPPVVSQGQRLSPGSY  
PTVIQQNATSQRAAKNGPPVSDQKEVLPAGVTIKQEQNLDQTYLDDELIDTHLSWIQNIL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	97.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.



[View online »](#)

<b>Note:</b>	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
<b>Storage:</b>	Store at -80°C.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_001129493</a>
<b>Locus ID:</b>	4773
<b>UniProt ID:</b>	<a href="#">Q13469</a>
<b>Cytogenetics:</b>	20q13.2
<b>RefSeq ORF:</b>	2703
<b>Synonyms:</b>	NFAT1; NFATP
<b>Summary:</b>	This gene is a member of the nuclear factor of activated T cells (NFAT) family. The product of this gene is a DNA-binding protein with a REL-homology region (RHR) and an NFAT-homology region (NHR). This protein is present in the cytosol and only translocates to the nucleus upon T cell receptor (TCR) stimulation, where it becomes a member of the nuclear factors of activated T cells transcription complex. This complex plays a central role in inducing gene transcription during the immune response. Alternate transcriptional splice variants encoding different isoforms have been characterized. [provided by RefSeq, Apr 2012]
<b>Protein Families:</b>	Druggable Genome, Transcription Factors
<b>Protein Pathways:</b>	Axon guidance, B cell receptor signaling pathway, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, T cell receptor signaling pathway, VEGF signaling pathway, Wnt signaling pathway

### Product images:



Coomassie blue staining of purified NFATC2 protein (Cat# TP326803). The protein was produced from HEK293T cells transfected with NFATC2 cDNA clone (Cat# [RC226803]) using MegaTran 2.0 (Cat# [TT210002]).