

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP326864

LMO2 (NM_001142316) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins	
Description:	Recombinant protein of human LIM domain only 2 (rhombotin-like 1) (LMO2), transcript variant 3, 20 μg	
Species:	Human	
Expression Host:	HEK293T	
Expression cDNA Clone or AA Sequence:	>RC226864 representing NM_001142316 Red=Cloning site Green=Tags(s)	
	MSSAIERKSLDPSEEPVDEVLQIPPSLLTCGGCQQNIGDRYFLKAIDQYWHEDCLSCDLCGCRLGEVGRR LYYKLGRKLCRRDYLRLFGQDGLCASCDKRIRAYEMTMRVKDKVYHLECFKCAACQKHFCVGDRYLLINS DIVCEQDIYEWTKINGMI	
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV	
Tag:	C-Myc/DDK	
Predicted MW:	18.2 kDa	
Concentration:	>0.05 µg/µL as determined by microplate BCA method	
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining	
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol	
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.	
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.	
Storage:	Store at -80°C.	
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.	
RefSeq:	<u>NP 001135788</u>	
Locus ID:	4005	
UniProt ID:	<u>P25791</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	LMO2 (NM_001142316) Human Recombinant Protein – TP326864	
Cytogenetics:	11p13	
RefSeq ORF:	474	
Synonyms:	LMO-2; RBTN2; RBTNL1; RHOM2; TTG2	
Summary:	LMO2 encodes a cysteine-rich, two LIM-domain protein that is required for yolk sac erythropoiesis. The LMO2 protein has a central and crucial role in hematopoietic development and is highly conserved. The LMO2 transcription start site is located approximately 25 kb downstream from the 11p13 T-cell translocation cluster (11p13 ttc), where a number T-cell acute lymphoblastic leukemia-specific translocations occur. Alternative splicing results in multiple transcript variants encoding different isoforms.[provided by RefSeq, Nov 2008]	
Protein Families	: Druggable Genome	

Product images:

116 -	-
66 -	-
45 -	-
35 -	- 11
25 -	-
18 -	-
14 -	-

Coomassie blue staining of purified LMO2 protein (Cat# TP326864). The protein was produced from HEK293T cells transfected with LMO2 cDNA clone (Cat# [RC226864]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US