

Product datasheet for TP327155

SPART (NM_001142294) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens spastic paraplegia 20 (Troyer syndrome) (SPG20), transcript variant 4, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC227155 protein sequence Red=Cloning site Green=Tags(s)
	<p>MEQEPQNGEPAEIKIIREAYKKAFLFVNKGLNTDELGQKEEAKNYYKQGIGHLLRGISISSKESEHTGTG WESARQMQQKMKETLQNVTRLEILEKGLATSLQNDLQEVPKLYPEFPPKDMCEKLPEPQSFSSAPQHA VNGNTSTPSAGAVAAPASLSLPSQSCPAEAPPAYTPQAAEGHYTVSYGTDSGEFSSVGEFVYRHSQPPP LETLGLDADELILIPNGVQIFFVNPAGEVSAPSYPGYLRIVRFLDNSLDTVLNRPPGFLQVCDWLPLYLP DRSPVLKCTAGAYMFPDTMLQAAGCFVGVVLSSELPEDDRELFEDLLRQMSDLRLQANWNRAEEENE PGRTRPSSDQLKEASGTDVKQLDQGNKDVRHKGKRGKRAKDTSSSEVNLSHIVPCEPVPEEKPKELHE WSEKVAHNILSGASWVSWGLVKGAIEITGKAIKQGASKLRERIQPEEKVPEVSPAVTKGLYIAKQATGGA AKV SQFLVDGVCTVANCVGKELAPHVKKHGSKLVPESLKKDKDGKSPLDGAMWVAASSVQGFSTVWQGLE CAA KCIVNNVSAETVQTVRYKYGYNAGEATHHAVDSAVNVGVTAYNINNIGIKAMVKKTATQTGHTLLE DYQI VDNSQRENQEGAANVNRGEKDEQTKVEKAKKKDK</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	72.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001135766
Locus ID:	23111
UniProt ID:	Q8N0X7 , A0A024RDV9
RefSeq Size:	4838
Cytogenetics:	13q13.3
RefSeq ORF:	1998
Synonyms:	SPG20; TAHCCP1
Summary:	This gene encodes a protein containing a MIT (Microtubule Interacting and Trafficking molecule) domain, and is implicated in regulating endosomal trafficking and mitochondria function. The protein localizes to mitochondria and partially co-localizes with microtubules. Stimulation with epidermal growth factor (EGF) results in protein translocation to the plasma membrane, and the protein functions in the degradation and intracellular trafficking of EGF receptor. Multiple alternatively spliced variants, encoding the same protein, have been identified. Mutations associated with this gene cause autosomal recessive spastic paraplegia 20 (Troyer syndrome). [provided by RefSeq, Nov 2008]

Product images:



Coomassie blue staining of purified SPART protein (Cat# TP327155). The protein was produced from HEK293T cells transfected with SPART cDNA clone (Cat# [RC227155]) using MegaTran 2.0 (Cat# [TT210002]).