

## Product datasheet for TP327811M

### DACH2 (NM\_001139515) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human dachshund homolog 2 (Drosophila) (DACH2), transcript variant 3, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC227811 representing NM_001139515 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)  MTRKQAVNSSRPGRPPKRSLGVLQENARLLTHAVPGLLSPGLITPTGITAAAMAEAMKLQKMKLMAMNTL QGNGSQNGTESEPDLLNSNTGGSESSWDKDKMQSPFAAPGPQHGIHAALAGQPGGGAPTLNPLQQNHL LTNRDLDPFMMMPHLLPVSLPPASVAMAMNQMNHLNNTIANMAAAAQIHSPLSRAGTSVIKERIPESPS APSLEENHRPGSQTSSHTSSSVSSSPSQMDHHLERMEEVPVQIPIMKSPLDKIQLTPGQALPAGFPGPFI FADSLSSVETLLTNIQGLLKVALDNARIQEKQIQEKKELRLELYREREIRENLERQLAVELQSRTTMQK RLKKEKTKRKLQEALFESKRREQVEQALKQATTSDSGLRMLKDTGIPDIEIENNGTPHDSAAMQGGNY YCLEMAQQLYSA  <b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-Myc/DDK
Predicted MW:	47 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP\\_001132987](#)

Locus ID: 117154

UniProt ID: [Q96NX9](#), [A8K311](#)


Cytogenetics: Xq21.2

RefSeq ORF: 1296

**Summary:** This gene is one of two genes which encode a protein similar to the *Drosophila* protein dachshund, a transcription factor involved in cell fate determination in the eye, limb and genital disc of the fly. The encoded protein contains two characteristic dachshund domains: an N-terminal domain responsible for DNA binding and a C-terminal domain responsible for protein-protein interactions. This gene is located on the X chromosome and is subject to inactivation by DNA methylation. The encoded protein may be involved in regulation of organogenesis and myogenesis, and may play a role in premature ovarian failure. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2008]

**Protein Families:** Transcription Factors

### Product images:



Coomassie blue staining of purified DACH2 protein (Cat# [TP327811]). The protein was produced from HEK293T cells transfected with DACH2 cDNA clone (Cat# [RC227811]) using MegaTran 2.0 (Cat# [TT210002]).