

Product datasheet for TP329337L

SETMAR (NM_006515) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human SET domain and mariner transposase fusion gene (SETMAR), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC229337 representing NM_006515 Red=Cloning site Green=Tags(s)

MFAEAAKTRPCGMAEFKEKPEAPTEQLDVACGQENLPVGAWPPGAAPPFQYTPDHVVGPGADIDPTQI
TFPGCICVKTPCLPGTCSCLRHHGENYDDNSCLRDIGSGGKYAEPVFECNLCRCSDHCRNRVWQKGLQFH
FQVFKTHKKGWGLRTLFIKGRFVCEYAGEVLGFSEVQRRHLQTKSDSNYIIAIREHVYNGQVMETFV
DPTYIGNIGRFLNHSCEPNLLMIPVRIDSMVPKLALFAAKDIVPEEELSYDYSGRYLNLTVSEDKERLDH
GKLRKPCYCGAKSCTAFLPFSSLYCPVEKSNISCGNEKEPSMCGSAPSVFPCKRLTLETMKMMLDKKQ
IRAIFFEFKMGRKAAETTRNINNAFGPGTANERTVQWWFKKFKGDESLEDEERSGRPSEVDNDQLRAI
IEADPLTTTREVAAELNVNHSTVVRHLKQIGKVKKLDKWWPHELTENQKNRRFEVSSSLLRNHNPEFLD
RIVTCDEKWLYDNRRRSQAQWLDQEEAPKHFPPKILHPKKVMVTIWWWSAAGLIHYSFLNPGETITSEKYA
QEIDEMNQKLQRLQALVNRKGPILLHDNARPHVAQPTLQKLNELGYEVLPHPPYSPDLLPTNYHVFKHL
NNFLQGKRFHNQQAENAFQEFVESQSTDFYATGINQLISRWQKCVDCNGSYFD


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	77.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_006506
Locus ID:	6419
UniProt ID:	Q53H47 , E7EN68
Cytogenetics:	3p26.1
RefSeq ORF:	2052
Synonyms:	Mar1; METNASE
Summary:	This gene encodes a fusion protein that contains an N-terminal histone-lysine N-methyltransferase domain and a C-terminal mariner transposase domain. The encoded protein binds DNA and functions in DNA repair activities including non-homologous end joining and double strand break repair. The SET domain portion of this protein specifically methylates histone H3 lysines 4 and 36. This gene exists as a fusion gene only in anthropoid primates, other organisms lack mariner transposase domain. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Jan 2013]
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Lysine degradation

Product images:

Coomassie blue staining of purified SETMAR protein (Cat# [TP329337]). The protein was produced from HEK293T cells transfected with SETMAR cDNA clone (Cat# [RC229337]) using MegaTran 2.0 (Cat# [TT210002]).