

Product datasheet for TP329900L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MICA (NM_001177519) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Homo sapiens MHC class I polypeptide-related sequence A

(MICA), transcript variant 1 (allele MICA*00801), 1 mg

Species: Human Expression Host: HEK293T

Expression cDNA Clone >RC229900 representing NM 001177519

or AA Sequence: Red=Cloning site Green=Tags(s)

MGLGPVFLLLAGIFPFAPPGAAAEPHSLRYNLTVLSWDGSVQSGFLAEVHLDGQPFLRYDRQKCRAKPQG QWAEDVLGNKTWDRETRDLTGNGKDLRMTLAHIKDQKEGLHSLQEIRVCEIHEDNSTRSSQHFYYDGELF LSQNLETEEWTVPQSSRAQTLAMNVRNFLKEDAMKTKTHYHAMHADCLQELRRYLESGVVLRRTVPPMVN VTRSEASEGNITVTCRASSFYPRNIILTWRQDGVSLSHDTQQWGDVLPDGNGTYQTWVATRICRGEEQRF

TCYMEHSGNHSTHPVPSGKVLVLQSHWQTFHVSAVAAGCCYFCYYYFLCPLL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 38.3

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: NULL or Add: Recombinant proteins was captured through anti-DDK affinity column followed

by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: <u>NP 001170990</u>

Locus ID: 100507436





UniProt ID: Q96QC4

Cytogenetics: 6p21.33

RefSeq ORF: 996

Synonyms: MIC-A; PERB11.1


Summary: This gene encodes the highly polymorphic major histocompatability complex class I chain-

related protein A. The protein product is expressed on the cell surface, although unlike canonical class I molecules it does not seem to associate with beta-2-microglobulin. It is a ligand for the NKG2-D type II integral membrane protein receptor. The protein functions as a stress-induced antigen that is broadly recognized by intestinal epithelial gamma delta T cells. Variations in this gene have been associated with susceptibility to psoriasis 1 and psoriatic arthritis, and the shedding of MICA-related antibodies and ligands is involved in the

progression from monoclonal gammopathy of undetermined significance to multiple myeloma. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jan

2014]

Product images:



Coomassie blue staining of purified MICA protein (Cat# [TP329900]). The protein was produced from HEK293T cells transfected with MICA cDNA clone (Cat# [RC229900]) using MegaTran 2.0 (Cat# [TT210002]).