

## **Product datasheet for TP330029**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ATXN10 (NM\_001167621) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human ataxin 10 (ATXN10), transcript variant 2, 20 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC230029 representing NM\_001167621

or AA Sequence: Red=Cloning site Green=Tags(s)

MAAPRPPPARLSGVMVPAPIQDLEALRALTALFKEQRNRNLDTIGVAVDLILLFRELRVEQESLLTAFRC GLQFLGNIASRNEDSQSIVWVHAFPELFLSCLNHPDKKIVAYSSMILFTSLNHERMKELEENLNIAIDVI DAYQKHPESEWPFLIITDLFLKSPELVQAMFPKLNNQERVTLLDLMIAKITSDEPLTKDDIPVFLRHAEL IASTFVDQCKTVLKLASEEPPDDEEALATIRLLDVLCEMTVNTELLGYLQVFPGLLERVIDLLRVIHVAG KETTNIFSNCGCVRAEGDISNVANGFKSHLIRLIGNLCYKNKDNQDKVNELDGIPLILDNCNISDSNPFL

TQWVIYAIRNLTEDNSQNQDLIAKMEEQGLADASLLKKVGFEVEKKGEKLILKSTRDTPKP

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK

Predicted MW: 46.7

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: NULL or Add: Recombinant proteins was captured through anti-DDK affinity column followed

by conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeg:** NP 001161093

**Locus ID:** 25814



ORIGENE

**UniProt ID:** Q9UBB4, A6NLC4

Cytogenetics: 22q13.31


RefSeq ORF: 1233

Synonyms: E46L; HUMEEP; SCA10

**Summary:** This gene encodes a protein that may function in neuron survival, neuron differentiation, and

neuritogenesis. These roles may be carried out via activation of the mitogen-activated protein kinase cascade. Expansion of an ATTCT repeat from 9-32 copies to 800-4500 copies in an intronic region of this locus has been associated with spinocerebellar ataxia, type 10. Alternatively spliced transcript variants have been described.[provided by RefSeq, Jul 2016]

## **Product images:**



Coomassie blue staining of purified ATXN10 protein (Cat# TP330029). The protein was produced from HEK293T cells transfected with ATXN10 cDNA clone (Cat# [RC230029]) using MegaTran 2.0 (Cat# [TT210002]).