

Product datasheet for **TP330258L**

PIP5K3 (PIKFYVE) (NM_001178000) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens phosphoinositide kinase, FYVE finger containing (PIKFYVE), transcript variant 4, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC230258 representing NM_001178000 Red =Cloning site Green =Tags(s)

MATDDKTSPTLDSANDLPRSPTSPSHLTHFKPLTPDQDEPPFKSAYSSFVNLFRFNKERAEGGQGEQQPL
SGSWTSPQLPSRTQSVRSPTYKKQLNEELQRRSSALDTRRKAEPTEFGGHDPRTAVQLRSLSTVLKRLKE
IMEGKSQDSDLKQYWMPDSQCKEYDCSEKFTTFRRRHHCRLCGQIFCSRCCNQEIPGKFMGYTGDLRAC
TYCRKIALSYAHSTDSNSIGEDLNALSACS SVSLDPSEPRTVPVGSRKASRNIFLEDDLAWQSLIHPDS
SNTPLSTRLVSVQEDAGKSPARNRSASITNLSLDRSGSPMVPSYETS VSPQANRTYVRTETTEDERKILL
DSVQLKDLWKKICHSSGMEFQDHRYLWRTHPNCIVGKELVNWLRNGHIATRAQAIAGQAMVDGRWLD
CVSHHDQLFRDEYALYRPLQSTEFSETPSPDSDSVNSVEGHSEPSWFKDIKFDDSDTEQIAEEGDDNLAN
SASPSKRTSVSSFQSTVSDSAASISLNVELDNVNFHIKKPSKYPHVPPHPADQKGRR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	62
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	NULL or Add: Recombinant proteins was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001171471
Locus ID:	200576
UniProt ID:	Q9Y2I7
Cytogenetics:	2q34
RefSeq ORF:	1644
Synonyms:	CFD; FAB1; HEL37; PIP5K; PIP5K3; ZFYVE29
Summary:	Phosphorylated derivatives of phosphatidylinositol (PtdIns) regulate cytoskeletal functions, membrane trafficking, and receptor signaling by recruiting protein complexes to cell- and endosomal-membranes. Humans have multiple PtdIns proteins that differ by the degree and position of phosphorylation of the inositol ring. This gene encodes an enzyme (PIKfyve; also known as phosphatidylinositol-3-phosphate 5-kinase type III or PIPKIII) that phosphorylates the D-5 position in PtdIns and phosphatidylinositol-3-phosphate (PtdIns3P) to make PtdIns5P and PtdIns(3,5)biphosphate. The D-5 position also can be phosphorylated by type I PtdIns4P-5-kinases (PIP5Ks) that are encoded by distinct genes and preferentially phosphorylate D-4 phosphorylated PtdIns. In contrast, PIKfyve preferentially phosphorylates D-3 phosphorylated PtdIns. In addition to being a lipid kinase, PIKfyve also has protein kinase activity. PIKfyve regulates endomembrane homeostasis and plays a role in the biogenesis of endosome carrier vesicles from early endosomes. Mutations in this gene cause corneal fleck dystrophy (CFD); an autosomal dominant disorder characterized by numerous small white flecks present in all layers of the corneal stroma. Histologically, these flecks appear to be keratocytes distended with lipid and mucopolysaccharide filled intracytoplasmic vacuoles. Alternative splicing results in multiple transcript variants encoding distinct isoforms.[provided by RefSeq, May 2010]
Protein Families:	Druggable Genome
Protein Pathways:	Endocytosis, Fc gamma R-mediated phagocytosis, Inositol phosphate metabolism, Metabolic pathways, Phosphatidylinositol signaling system, Regulation of actin cytoskeleton

Product images:



Coomassie blue staining of purified PIKFYVE protein (Cat# [TP330258]). The protein was produced from HEK293T cells transfected with PIKFYVE cDNA clone (Cat# [RC230258]) using MegaTran 2.0 (Cat# [TT210002]).