

Product datasheet for **TP500390**

Rbx1 (NM_019712) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Mouse ring-box 1 (Rbx1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >MR200390 protein sequence
Red=Cloning site **Green**=Tags(s)

MAAAMDVDTPSGTNSGAGKKRFEVKKWNAVALWAWDIVDNCIAICRNHIMDLICIEQANQASATSEECTV
AWGVCNHAHFHFCISRWLKTRQVCPLDNREWEFQKYGH

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-MYC/DDK

Predicted MW: 12.3 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_062686](#)

Locus ID: 56438

UniProt ID: [P62878](#)

RefSeq Size: 1655

Cytogenetics: 15 E1

RefSeq ORF: 327



[View online »](#)

Synonyms: 1500002P15Rik; AA517855; ROC1

Summary: E3 ubiquitin ligase component of multiple cullin-RING-based E3 ubiquitin-protein ligase (CRLs) complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins, including proteins involved in cell cycle progression, signal transduction, transcription and transcription-coupled nucleotide excision repair (PubMed:22118460). CRLs complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins, ARIH1 mediating addition of the first ubiquitin on CRLs targets (By similarity). The functional specificity of the E3 ubiquitin-protein ligase complexes depends on the variable substrate recognition components (By similarity). As a component of the CSA complex promotes the ubiquitination of ERCC6 resulting in proteasomal degradation (By similarity). Through the RING-type zinc finger, seems to recruit the E2 ubiquitination enzyme, like CDC34, to the complex and brings it into close proximity to the substrate (By similarity). Probably also stimulates CDC34 autoubiquitination (By similarity). May be required for histone H3 and histone H4 ubiquitination in response to ultraviolet and for subsequent DNA repair (By similarity). Promotes the neddylation of CUL1, CUL2, CUL4 and CUL4 via its interaction with UBE2M (By similarity). Involved in the ubiquitination of KEAP1, ENC1 and KLHL41 (By similarity). In concert with ATF2 and CUL3, promotes degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM (By similarity).[UniProtKB/Swiss-Prot Function]