

Product datasheet for TP500743

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Med31 (NM 026068) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Mouse mediator complex subunit 31 (Med31), with C-terminal

MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse

Expression Host: HEK293T

Expression cDNA Clone >MR200743 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAAAVAMETDDAGNRLRFQLELEFVQCLANPNYLNFLAQRGYFKDKAFVNYLKYLLYWKEPEYAKYLKYP

QCLHMLELLQYEHFRKELVNAQCAKFIDEQQILHWQHYSRKRVRLQQALAEQQQQNNTAGK

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-MYC/DDK

Predicted MW: 15.8 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C after receiving vials.

11 B4

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 080344

 Locus ID:
 67279

 UniProt ID:
 Q9CXU1

 RefSeq Size:
 1179

RefSeq ORF: 396

Cytogenetics:





Med31 (NM_026068) Mouse Recombinant Protein - TP500743

Synonyms: 3110004H13Rik; CGI-125; l11Jus15

Summary: Component of the Mediator complex, a coactivator involved in the regulated transcription of

nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (By similarity).

[UniProtKB/Swiss-Prot Function]