

## **Product datasheet for TP500914**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Med9 (NM\_138675) Mouse Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse mediator complex subunit 9 (Med9), with C-terminal

MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse

**Expression Host:** HEK293T

**Expression cDNA Clone** >MR200914 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MASSGVAGGRQAEDTLQPPPELLPESKPPPPPQPLPVAALPPPAAPRPQSPAGAKEENYSFLPLVHNVIK CMDKDSPDLHQDLNALKTKFQELRKLIGTMPGIHVSPEQQQQQLHSLREQVRTKNELLQKYKSLCMFEIP

KD

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-MYC/DDK

Predicted MW: 15.7 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

**Storage:** Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 619616

 Locus ID:
 192191

 UniProt ID:
 Q8VCS6

 RefSeq Size:
 3013

Cytogenetics: 11 B1.3





## Med9 (NM\_138675) Mouse Recombinant Protein - TP500914

RefSeq ORF: 429

Synonyms: BC019367; Med25

**Summary:** Component of the Mediator complex, a coactivator involved in the regulated transcription of

nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (By similarity).

[UniProtKB/Swiss-Prot Function]