

Product datasheet for TP500965

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Trfp (BC060122) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Mouse Trf (TATA binding protein-related factor)-proximal

protein homolog (Drosophila) (cDNA clone MGC:63299, with C-terminal MYC/DDK tag, expressed

in HEK293T cells, 20ug

Species: Mouse

Expression Host: HEK293T

Expression cDNA Clone >MR200965 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MGVTCVSQMPVAEGKSLQQTVELLTKKLEMLGAEKQGTFCVDCETYHTAASTLGSQGQAGKLMYVMHNSE YPLSCFALFENGPCLIADTNFDVLMVKLKGFFQSAKASKIETRGTRYQYCDFLVKVGTVTMGPSARGISV

EVRPW

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-MYC/DDK

Predicted MW: 15.9 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

 Locus ID:
 56771

 UniProt ID:
 Q9R0X0

 RefSeq Size:
 5774

 Cytogenetics:
 17 C





Trfp (BC060122) Mouse Recombinant Protein - TP500965

RefSeq ORF: 435

Synonyms: 1110011005Rik; 2410115I17Rik; AU018348; Trfp

Summary: Component of the Mediator complex, a coactivator involved in the regulated transcription of

nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (By similarity).[UniProtKB/Swiss-Prot

Function]