

Product datasheet for TP501030

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Crcp (NM_007761) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Mouse calcitonin gene-related peptide-receptor component

protein (Crcp), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse

Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

>MR201030 protein sequence Red=Cloning site Green=Tags(s)

MEVKDANAALLSNYEVFQLLTDLKEQRKESGKNKHSAGQQNLNAITYETLKYISKTPCRNQSPAIVQEFL TAMKSHKLTKAEKLQLLNHRPMTAVEIQLMVEESEERLTEEQIEALLHTVTSILPAGPEDEQSKSTSNDV

AMEEEEPA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-MYC/DDK

Predicted MW: 16.7 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 031787

 Locus ID:
 12909

 UniProt ID:
 035427

 RefSeq Size:
 1474

Cytogenetics: 5 G1.3





Crcp (NM_007761) Mouse Recombinant Protein - TP501030

RefSeq ORF: 447

Synonyms: AL022669

Summary: DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four

ribonucleoside triphosphates as substrates. Specific peripheric component of RNA

polymerase III which synthesizes small RNAs, such as 5S rRNA and tRNAs. Plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts

induce type I interferon and NF- Kappa-B through the RIG-I pathway (By similarity).

[UniProtKB/Swiss-Prot Function]