OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for TP501073

Wtap (NM_175394) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse Wilms tumour 1-associating protein (Wtap), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR201073 protein sequence Red=Cloning site Green=Tags(s) MTNEEPLPKKVRLSETDFKVMARDELILRWKQYEAYVQALEGKYTDLNSNDVTGLRESEEKLKQQQQESA RRENILVMRLATKEQEMQECTTQIQYLKQVQQPSVAQLRSTMVDPAINLFFLKMKGELEQTKDKLEQAQ N ELSAWKFTPDR TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-MYC/DDK
Predicted MW:	17.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_780603
Locus ID:	60532
UniProt ID:	Q9ER69
RefSeq Size:	4967

View online »

Cytogenetics:	17 A1
RefSeq ORF:	453
Synonyms:	2810408K05Rik; 9430038B09Rik
Summary:	Associated component of the WMM complex, a complex that mediates N6-methyladenosine (m6A) methylation of RNAs, a modification that plays a role in the efficiency of mRNA splicing and RNA processing (PubMed:29535189, PubMed:29547716). Acts as a key regulator of m6A methylation by promoting m6A methylation of mRNAs at the 3' UTR (PubMed:29547716). Required for accumulation of METTL3 and METTL14 to nuclear speckle (By similarity). Acts as a mRNA splicing regulator (By similarity). Regulates G2/M cell-cycle transition by binding to the 3' UTR of CCNA2, which enhances its stability (By similarity). Impairs WT1 DNA-binding ability and inhibits expression of WT1 target genes (By similarity).[UniProtKB/Swiss-Prot Function]