

## Product datasheet for TP501830

### Cdc42 (NM\_009861) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse cell division cycle 42 (Cdc42), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR201830 representing NM_009861 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)  MQTIKCWWGDGAVGKTCLLISYTTNKFSEYVPTVFDNYAVTMIGGEPYTLGLFDTAGQEDYDGLRPL SYPQTDVFLVCFVSPSSFENVKEKWVPEITHHCPKTPFLLVGTQIDLRDDPSTIEKLAKNKQKPITPE TAEKLARDLKAVKYVECSALTQKGLKNVFDGAILAALEPPEPKSRRCVLL  <b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-MYC/DDK
Predicted MW:	21.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u><a href="#">NP_033991</a></u>
Locus ID:	12540
UniProt ID:	<u><a href="#">P60766</a></u>
RefSeq Size:	2063
Cytogenetics:	4 69.83 cM


[View online »](#)

**RefSeq ORF:** 573

**Synonyms:** AI747189; AU018915

**Summary:** Plasma membrane-associated small GTPase which cycles between an active GTP-bound and an inactive GDP-bound state (PubMed:24352656). In its active state binds to a variety of effector proteins to regulate cellular responses. Involved in epithelial cell polarization processes. Regulates the bipolar attachment of spindle microtubules to kinetochores before chromosome congression in metaphase. Regulates cell migration (By similarity). In neurons, plays a role in the extension and maintenance of the formation of filopodia, thin and actin-rich surface projections. Required for DOCK10-mediated spine formation in Purkinje cells and hippocampal neurons (PubMed:25851601). Facilitates filopodia formation upon DOCK11-activation (PubMed:22494997). Upon activation by CaMKII, modulates dendritic spine structural plasticity by relaying CaMKII transient activation to synapse-specific, long-term signaling (By similarity). Also plays a role in phagocytosis through organization of the F-actin cytoskeleton associated with forming phagocytic cups (By similarity).[UniProtKB/Swiss-Prot Function]