

## **Product datasheet for TP502131**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Polr3h (NM\_030229) Mouse Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

Description: Purified recombinant protein of Mouse polymerase (RNA) III (DNA directed) polypeptide H

(Polr3h), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse

**Expression Host:** HEK293T

**Expression cDNA Clone** >MR202131 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MFVLVEMVDTVRIPPWQFERKLNDSIAEELNKKLANKVVYNVGLCICLFDITKLEDAYVFPGDGASHTKV HFRYVVFHPFLDEILIGKIKGCSPEGVHVSLGFFDDILIPPESLQQPAKFDEAEQVWVWEYETEEGAHDL YMDTGEEIRFRVVDESFVDTSPTGPSSAEAASSSEELPKKEAPYTLVGSISEPGLGLLSWWTSN

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-MYC/DDK

Predicted MW: 22.9 kDa

**Concentration:**  $>0.05 \mu g/\mu L$  as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

**Storage:** Store at -80°C after receiving vials.

15 E1

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** <u>NP 084505</u>

 Locus ID:
 78929

 UniProt ID:
 Q9D2C6

 RefSeq Size:
 2514

Cytogenetics:





## Polr3h (NM\_030229) Mouse Recombinant Protein - TP502131

RefSeq ORF: 615

**Synonyms:** 5031409G22Rik; RPC8

Summary: DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four

ribonucleoside triphosphates as substrates. Specific peripheric component of RNA

polymerase III which synthesizes small RNAs, such as 5S rRNA and tRNA. Plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts

induce type I interferon and NF- Kappa-B through the RIG-I pathway (By similarity).

[UniProtKB/Swiss-Prot Function]