

## Product datasheet for **TP502493**

### Chmp5 (NM\_029814) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse charged multivesicular body protein 5 (Chmp5), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR202493 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	<p>MNRRFFGKAKPKAPPPSLTDCIGTVDSRAESIDKKISRLDAELVKYKDQIKKMREGPAKNMVKQKALRVLK QKRMYEQQRDNLAQSFNMEQANYTIQSLKDKTTVDAMKLGVKEMKKAYKEVKIDQIEDLQDQLEDMMME DANEIQEALGRSYGTPELDEDDLEAELDALGDELLADESSYLDEAASAPAIPEGVPTDTKNKDGVLVDE FGLPQIPAS</p> <p><b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b></p>
Tag:	C-MYC/DDK
Predicted MW:	24.6 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_084090</a>
Locus ID:	76959
UniProt ID:	<a href="#">Q9D7S9</a>
RefSeq Size:	1372



[View online »](#)

**Cytogenetics:** 4 A5

**RefSeq ORF:** 660

**Synonyms:** 2210412K09Rik; AW545668

**Summary:** Probable peripherally associated component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I, -II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis. ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4 (By similarity). [UniProtKB/Swiss-Prot Function]