

Product datasheet for TP502602

Chmp3 (NM_025783) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse charged multivesicular body protein 3 (Chmp3), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR202602 protein sequence Red =Cloning site Green =Tags(s)
	MGLFGKTQEPPKELVNEWSLKIRKEMRWDRQIRDIQREEEKVKRSVKDAAKKGQKEVCVVLAKEMIRS RKAVSKLYASKAHMNSVLMGMKNQLAVLRVAGSLQKSTEVMMKAMQSLVKIPEIQATMRELSKEMMKAGI EEMLEDTFESMDDQEEMEEAAEMEIDRILFEITAGALGKAPSKVTDALPEPEPAGAMAASEEGEEEEDEE DLEAMQSRLATLRS
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-MYC/DDK
Predicted MW:	25.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_080059
Locus ID:	66700
UniProt ID:	Q9CQ10



[View online »](#)

RefSeq Size:	2592
Cytogenetics:	6 32.17 cM
RefSeq ORF:	672
Synonyms:	25.1; 4921505F14Rik; 9130011K15Rik; CGI-49; D6Ertd286e; NEDF; Vps24
Summary:	<p>Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I, -II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis. ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. Selectively binds to phosphatidylinositol 3,5-bisphosphate PtdIns(3,5)P2 and PtdIns(3,4)P2 in preference to other phosphoinositides tested. Involved in late stages of cytokinesis. Plays a role in endosomal sorting/trafficking of EGF receptor (By similarity). [UniProtKB/Swiss-Prot Function]</p>