

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP502607

Chmp4b (NM_029362) Mouse Recombinant Protein

Product data:

Recombinant Proteins
Purified recombinant protein of Mouse charged multivesicular body protein 4B (Chmp4b), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Mouse
HEK293T
>MR202607 protein sequence Red=Cloning site Green=Tags(s)
MSVFGKLFGAGGGKAGKGGPTPQEAIQRLRDTEEMLSKKQEFLEKKIEQELTAAKKHGTKNKRAALQALK RKKRYEKQLAQIDGTLSTIEFQREALENANTNTEVLKNMGYAAKAMKAAHDNMDIDKVDELMQDIADQQ E
LAEEISTAISKPVGFGEEFDEDELMAELEELEQEELDKNLLEISGPETVPLPNVPSVALPSKPAKKKEEE DDDMKELENWAGSM
TRTRPLEQKLISEEDLAANDILDYKDDDDKV
C-MYC/DDK
24.9 kDa
>0.05 µg/µL as determined by microplate BCA method
> 80% as determined by SDS-PAGE and Coomassie blue staining
25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Store at -80°C after receiving vials.
Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<u>NP 083638</u>
75608
<u>Q9D8B3</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Chmp4b (NM_029362) Mouse Recombinant Protein – TP502607
RefSeq Size:	1607
Cytogenetics:	2 H1
RefSeq ORF:	672
Synonyms:	2010012F05Rik; C76846; Snf7-2
Summary:	Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I,-II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis. Together with SPAST, the ESCRT-III complex promotes nuclear envelope sealing and mitotic spindle disassembly during late anaphase. Plays a role in the endosomal sorting pathway. ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. When overexpressed, membrane-assembled circular arrays of CHMP4B filaments can promote or stabilize negative curvature and outward budding. CHMP4A/B/C are required for the exosomal release of SDCBP, CD63 and syndecan.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US