

Product datasheet for **TP502874**

Cd81 (NM_133655) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse CD81 antigen (Cd81), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR202874 protein sequence Red =Cloning site Green =Tags(s) MGVEGCTKCIKYLLFVFNFWLGGVILGVALWLRHDPQTTSLLYLELGNKPAPNTFYVGIYILIAVGA VMMFVGFLGCGAIQESQCLLGTTFTCLVILFACEVAAGIWGFVNKDQIAKDVQKFYDQALQQAVMDDD A NNAKAVVKTFFHETLNCCGSNALTTLTTLRNSLCPSGGNLTPLLQQDCHQKIDELFSGKLYLIGIAAI WVAVIMIFEMILSMVLCCGIRNSSVY TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-MYC/DDK
Predicted MW:	25.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_598416</u>
Locus ID:	12520
UniProt ID:	<u>P35762</u>


[View online »](#)

RefSeq Size:	1534
Cytogenetics:	7 88.1 cM
RefSeq ORF:	708
Synonyms:	Tapa-1; Tapa1; Tspan28
Summary:	<p>Structural component of specialized membrane microdomains known as tetraspanin-enriched microdomains (TERMs), which act as platforms for receptor clustering and signaling. Essential for trafficking and compartmentalization of CD19 receptor on the cell surface of activated B cells (PubMed:23499492). Upon initial encounter with a microbial pathogen, enables the assembly of CD19-CR2 and B cell receptor complexes at signaling TERMS, lowering the threshold dose of antigen required to trigger B cell clonal expansion and humoral immune response (By similarity). In T cells, associates with CD4 or CD8 coreceptors and defines the maturation state of antigen-induced synapses with B cells (By similarity). Facilitates localization of CD3 in these immune synapses, required for costimulation and sustained activation of T cells, preferentially triggering T helper type 2 immune response (PubMed:11046035). Can act both as positive and negative regulator of homotypic or heterotypic cell-cell fusion processes. In myoblasts, associates with another tetraspanin CD9 in complex with PTGFRN and inhibits myotube fusion during muscle regeneration (PubMed:23575678). In macrophages, associates with CD9 and beta-1 and beta-2 integrins, and prevents macrophage fusion into multinucleated giant cells specialized in ingesting complement-opsonized large particles. Also prevents the fusion between mononuclear cell progenitors into osteoclasts in charge of bone resorption. Positively regulates sperm-egg fusion and may be involved in the acrosome reaction (PubMed:16380109, PubMed:17290409). Regulates protein trafficking in intracellular compartments. In T cells, associates with dNTPase SAMHD1 and defines its subcellular location, enabling its degradation by the proteasome and thereby controlling intracellular dNTP levels (By similarity). Also regulates integrin-dependent migration of macrophages, particularly relevant for inflammatory response in the lung (PubMed:18662991).[UniProtKB/Swiss-Prot Function]</p>