

Product datasheet for TP503141

Psma7 (NM_011969) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse proteasome (prosome, macropain) subunit, alpha type 7 (Psma7), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR203141 protein sequence Red =Cloning site Green =Tags(s)
	MSYDRAITVFSPDGHFLFQVEYAQEA VKKGSTAVGVRGKDIWLVGVEKKSVAKLQDERTVRKICALDDNVC MAFAGLTADARIVINRARVEQCQSHRLTVEDPVTVEYITRYIASLKQRYTQSNRRPFGISALIVGFDFDG TPRLYQTDPSGTYHAWKANAIGRGAKSVREFLEKNYTDDAIETDDLTIKLVIKALLEVVQSGGKNIELAV MRRDQPLKILNPEEIEKYVAEIEKEKEENEKQKAS
	TR TRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-MYC/DDK
Predicted MW:	27.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_036099
Locus ID:	26444
UniProt ID:	Q9Z2U0 , Q542H2 , Q3UIT9
RefSeq Size:	952



[View online »](#)

Cytogenetics: 2 H4

RefSeq ORF: 747

Synonyms: C6-I

Summary: Component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP-dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex). [UniProtKB/Swiss-Prot Function]