

## Product datasheet for **TP503277**

### **Tgfb2 (BC011055) Mouse Recombinant Protein**

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse transforming growth factor, beta 2 (cDNA clone MGC:7998 IMAGE:3585774), complete cds, with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR203277 protein sequence <span style="color: red;">Red</span> =Cloning site <span style="color: green;">Green</span> =Tags(s)  MFRVEEATHVQILKSKDLTSPTQRYIDSKVVKTRAEGEWLSFDVTDVAVQEWLHHKDRNLGFKISLHCPCC TFVPSNNYIIPNKSEELARFAGIDGTSTYASGDQKTIKSTRKKTSGKTPHLLMLLPSYRLESQQSSRR KKRALDAAYCFRNVQDNCCLRPYIDFKRDLGWKWIHEPKGYNANFCAGACPYLWSSDTQHTKVLSLYN T INPEASASPCCVSQDLEPLTILYYIGNTPKIEQLSNMIVKSKCKS  <span style="color: red;">TR</span> <span style="color: green;">TRPLEQKLISEEDLAANDILDYKDDDDKV</span>
Tag:	C-MYC/DDK
Predicted MW:	29.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
Locus ID:	21808
UniProt ID:	<u><a href="#">P27090</a></u>
RefSeq Size:	2047


[View online »](#)

<b>Cytogenetics:</b>	1 89.95 cM
<b>RefSeq ORF:</b>	765
<b>Synonyms:</b>	BB105277; Tgf-beta2; Tgfb-2
<b>Summary:</b>	<p>This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGF-beta family members. Mice lacking a functional copy of this gene display developmental defects in multiple organs and perinatal lethality. Heterozygous mutant mice exhibit aortic root aneurysm. This gene encodes multiple isoforms that may undergo similar proteolytic processing. [provided by RefSeq, Aug 2016]</p>